### For Research Use Only

# **HLA-G Monoclonal antibody**

Catalog Number:66447-1-lg 25 Publications



**Purification Method:** 

Protein A purification

Recommended Dilutions:

WB 1:2000-1:20000 IHC 1:50-1:500

CloneNo.:

1E5A10

**Basic Information** 

Catalog Number: 66447-1-lg Concentration:

1300 µg/ml Source: Mouse Isotype: lgG2b

Immunogen Catalog Number:

AG10839

GenBank Accession Number:

BC021708 GeneID (NCBI): 3135 **UNIPROT ID:** 

Full Name: major histocompatibility complex, class I. G

Calculated MW: 338 aa, 38 kDa

P17693

Observed MW: 33-45 kDa

**Applications** 

**Tested Applications:** WB, IHC, ELISA

**Cited Applications:** WB, IHC, IF

Species Specificity:

human **Cited Species:** 

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: human placenta tissue, IHC: human placenta tissue,

## **Background Information**

 $Human\,major\,histocompatibility\,complex\,(MHC)\,antigens,\,also\,referred\,to\,as\,human\,leukocyte\,antigens\,(HLA),\,are$ encoded by genes located on the short arm of chromosome 6 (6p21.3). HLA-G is a non-classical MHC class I molecule with multiple immunoregulatory properties. HLA-G exhibits a restricted pattern of expression that includes placental extravillous trophoblasts at the maternal-fetal interface, where it abolishes maternal immune cell activity against fetus and establishes immune tolerance. Aberrant expression of HLA-G has been found in a variety of human neoplastic diseases. It plays an important role in the escape of tumor cells from immunosurveillance.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Liling Xiong	34605151	Aging Cell	IHC,IF
Manni Sun	33205477	FASEB J	IHC
Jingmei Ma	36420138	Front Cell Dev Biol	IHC,IF

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**



human placenta tissue were subjected to SDS PAGE followed by western blot with 66447-1-1g (HLA-G Antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 66447-1-1g (HLA-G Antibody) at dilution of 1:400 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human placenta tissue slide using 66447-1-lg (HLA-G Antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).