

For Research Use Only

STAT5A Monoclonal antibody

Catalog Number: 66459-1-Ig **1 Publications**



Basic Information

Catalog Number: 66459-1-Ig	GenBank Accession Number: BC027036	Purification Method: Protein A purification
Size: 2000 µg/ml	GeneID (NCBI): 6776	CloneNo.: 1F10F6
Source: Mouse	UNIPROT ID: P42229	Recommended Dilutions: WB 1:2000-1:16000 IF 1:200-1:800
Isotype: IgG2a	Full Name: signal transducer and activator of transcription 5A	
Immunogen Catalog Number: AG19546	Calculated MW: 794 aa, 92 kDa	
	Observed MW: 95 kDa	

Applications

Tested Applications: IF/ICC, WB, ELISA	Positive Controls:
Cited Applications: WB	WB : RAW 264.7 cells, U-937 cells, rat spleen tissue, pig spleen tissue
Species Specificity: human, mouse, pig, rat	IF : HepG2 cells,
Cited Species: human	

Background Information

STAT5A, also named as STAT5, belongs to the transcription factor STAT family. In response to cytokines and growth factors, STAT family members are phosphorylated by the receptor associated kinases, and then form homo- or heterodimers that translocate to the cell nucleus where they act as transcription activators. STAT5A is activated by, and mediates the responses of many cell ligands, such as IL2, IL3, IL7 GM-CSF, erythropoietin, thrombopoietin, and different GHs. Activation of STAT5A in myeloma and lymphoma associated with a TEL/JAK2 gene fusion is independent of cell stimulus and has been shown to be essential for the tumorigenesis. The mouse counterpart of human STAT5A is found to induce the expression of BCL2L1/BCL-X(L), which suggests the antiapoptotic function of STAT5A in cells.

Notable Publications

Author	Pubmed ID	Journal	Application
Xinyi Jiang	36927693	J Clin Invest	WB


Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66459-1-Ig (STAT5A antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using STAT5A antibody (66459-1-Ig, Clone: 1F10F6) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).