For Research Use Only

CRABP2 Monoclonal antibody, PBS Only **Foroteintech**

Catalog Number:66468-1-PBS

Featured Product




This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Antibodies | ELISA kits | Proteins

www.ptglab.com


Selected Validation Data




T-47D cells were subjected to SDS PAGE followed by western blot with 66468-1-1g (CRABP2 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66468-1-PBS in a different storage buffer formulation.




WB result of CRABP2 antibody (66468-1-Ig; 1:500; incubated at room temperature for 1.5 hours) with sh-Control and sh-CRABP2 transfected HEK-293 cells. This data was developed using the same antibody clone with 66468-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed human skin cancer tissue using CRABP2 antibody (66468-1-1g, Clone: 1A5F3) at dilution of 1:100 and Coralite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66468-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 66468-1-Ig (CRABP2 antibody) at dilution of 1:500 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66468-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human ovary tumor tissue slide using 66468-1-Ig (CRABP2 antibody) at dilution of 1:500 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66468-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed MCF-7 cells using CRABP2 antibody (66468-1-Ig, Clone: 1A5F3) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1). This data was developed using the same antibody clone with 66468-1-PBS in a different storage buffer formulation.



1X10⁶ MCF-7 cells were intracellularly stained with 0.4 ug Anti-Human CRABP2 (66468-1-Ig, Clone:1A5F3) and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG1 Isotype Control (MOPC-21) (65124-1-Ig, Clone: MOPC-21) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 66468-1-PBS in a

