For Research Use Only

EZH2 Monoclonal antibody

Catalog Number:66476-1-lg Featured Product

9 Publications

BC010858

2146

GeneID (NCBI):

Calculated MW: 751 aa, 86 kDa

GenBank Accession Number:



Basic Information

Catalog Number: 66476-1-lg Concentration: 1000 ug/ml

UNIPROT ID: Source: Mouse Q15910 Full Name: Isotype: lgG1 enhancer of zeste homolog 2

(Drosophila)

Immunogen Catalog Number:

AG16789

Observed MW: 90-102 kDa

Purification Method:

Protein G purification

CloneNo.: 1F10A12

Recommended Dilutions:

WB: 1:5000-1:20000

FC (Intra): 0.20 ug per 10^6 cells in a

100 µl suspension

Applications

Tested Applications: WB, FC (Intra), ELISA **Cited Applications:** WB, IHC, IF Species Specificity:

human, mouse, rat Cited Species: human, mouse

Positive Controls:

WB: HEK-293 cells, A549 cells, Jurkat cells, ROS1728 cells, NIH/3T3 cells, 4T1 cells, A431 cells, PC-3 cells,

DU145 cells

FC (Intra): HepG2 cells,

Background Information

EZH2 (enhancer of zeste homologue 2, also known as KMT6) is a member of Polycomb group (PcG) family and encodes a histone methyl transferase that has an essential role in promoting histone H3 lysine 27 trimethylation (H3K27me3) and epigenetic gene silencing. EZH2 is important for cell proliferation and inhibition of cell differentiation, and is implicated in cancer progression. Overexpression of EZH2 is a marker of advanced and metastatic disease in many solid tumors, including prostate and breast cancer. This antibody detected EZH2 protein as a single band with a molecular weight (MW) of 91-100 kDa in multiple cell lines. The phosphorylation may result in the higher molecular weight (calculated MW as 80-86 kDa). (20935635, 21367748)

Notable Publications

Author	Pubmed ID	Journal	Application
Nicholas Marano	36274837	Front Cell Dev Biol	WB
Longyang Jin	30282996	Cell Death Dis	WB
Ji Wang	36479622	Cancer Med	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various cell lines were subjected to SDS PAGE followed by western blot with 66476-1-1g (EZH2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



WB result of EZH2 antibody (66476-1-lg; 1:30000; incubated at room temperature for 1.5 hours) with sh-Control and sh-EZH2 transfected HEK-293 cells.



1X10^6 HepG2 cells were intracellularly stained with 0.2 ug Anti-Human EZH2 (66476-1-1g, Clone:1F10A12) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (green), and 0.2 ug Mouse IgG1 Isotype Control (66360-1-1g, Clone: T1F8D3F10) (black). Cells were fixed with 4% PFA and permeabilized with 0.1% TritonX-100.



Various lysates were subjected to SDS PAGE followed by western blot with 66476-1-1g (EZH2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.