For Research Use Only

EZH2 Monoclonal antibody, PBS Only

Catalog Number:66476-1-PBS Featured Product



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method:

66476-1-PBS

GeneID (NCBI):

Protein A purification

Size: 1 mg/ml

2146 **UNIPROT ID:**

BC010858

CloneNo.: 1F10A12

Source: Mouse

Q15910 Full Name:

Isotype: lgG1

enhancer of zeste homolog 2 (Drosophila)

Immunogen Catalog Number:

Calculated MW:

AG16789

751 aa, 86 kDa Observed MW: 90-102 kDa

Applications

Tested Applications:

WB, FC (Intra), Indirect ELISA

Species Specificity:

human, mouse, rat

Background Information

EZH2 (enhancer of zeste homologue 2, also known as KMT6) is a member of Polycomb group (PcG) family and encodes a histone methyl transferase that has an essential role in promoting histone H3 lysine 27 trimethylation (H3K27me3) and epigenetic gene silencing. EZH2 is important for cell proliferation and inhibition of cell differentiation, and is implicated in cancer progression. Overexpression of EZH2 is a marker of advanced and metastatic disease in many solid tumors, including prostate and breast cancer. This antibody detected EZH2 protein as a single band with a molecular weight (MW) of 91-100 kDa in multiple cell lines. The phosphorylation may result in the higher molecular weight (calculated MW as 80-86 kDa). (20935635, 21367748)

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

PBS Only


Selected Validation Data




Various cell lines were subjected to SDS PAGE followed by western blot with 66476-1-lg (EZH2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66476-1-PBS in a different storage buffer formulation.



WB result of EZH2 antibody (66476-1-Ig; 1:30000; incubated at room temperature for 1.5 hours) with sh-Control and sh-EZH2 transfected HEK-293 cells. This data was developed using the same antibody clone with 66476-1-PBS in a different storage buffer formulation.



1X10^6 HepG2 cells were intracellularly stained with 0.2 ug Anti-Human EZH2 (66476-1-lg, Clone:1F10A12) and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (green), and 0.2 ug Mouse IgG1 Isotype Control (66360-1-lg, Clone: T1F8D3F10) (black). Cells were fixed with 4% PFA and permeabilized with 0.1% TritonX-100. This data was developed using the same antibody clone with 66476-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 66476-1-lg (EZH2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66476-1-PBS in a different storage buffer formulation.