For Research Use Only

IgM Monoclonal antibody

Catalog Number:66484-1-lg 3 Publications



Basic Information

Catalog Number: GenBank Accession Number: 66484-1-lg BC009851

Protein G purification

Protein G purification

66484-1-lg Size: 1500 μg/ml

Source:

 GeneID (NCBI):
 CloneNo.:

 3507
 2D10B10

 Full Name:
 Recomme

Observed MW:

75 kDa

Recommended Dilutions: u WB 1:5000-1:20000

Mouse Isotype: IgG1 immunoglobulin heavy constant mu WB 1:5000-1:20000
Calculated MW: IHC 1:2000-1:6000
69 kDa IF 1:200-1:800

Immunogen Catalog Number: AG1459

Applications

Tested Applications:

IF/ICC,IF-P, IHC, WB, ELISA

Cited Applications: IF, IHC, WB

Species Specificity:

Human
Cited Species:

human

Positive Controls:

WB: Human plasma tissue,

IHC: human tonsillitis tissue, human lymphoma tissue

IF: human tonsillitis tissue, human appendicitis tissue

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

IGHM is the constant region of heavy chain of IgM. IgM is the first immunoglobulin expressed during B cell development. IgM antibodies play an important role in primary immune response involved in early recognition of external invaders like bacteria and viruses, cellular waste and modified self, as well as in recognition and elimination of precancerous and cancerous lesions. The $\,\mu$ heavy chain disease is a rare lymphoproliferative disorder. Human immunoglobulin heavy chain locus translocations are associated with leukaemias and lymphomas, including multiple myeloma, mantle cell lymphoma, Burkitt's lymphoma and diffuse large B cell lymphoma. This antibody detects the heavy chain of human IgM (~75 kDa).

Notable Publications

Author	Pubmed ID	Journal	Application
Han Liu	36106364	Adv Sci (Weinh)	IF
Anne Kathrin Wessig	35296708	Sci Rep	WB
Haishi Zheng	37933335	J Inflamm Res	IHC

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data




0.1 $\,\mu$ L human plasma was subjected to SDS PAGE followed by western blot with 66484-1-1g (IGHM antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using human IgM antibody (66484-1-Ig, Clone: 2D10B10) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66484-1-1g (IGHM antibody) at dilution of 1:1000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66484-1-Ig (IGHM antibody) at dilution of 1:1000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66484-1-1g (IGHM antibody) at dilution of 1:1000 (under 4x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human appendicitis tissue using human IgM antibody (66484-1-Ig, Clone: 2D10B10) at dilution of 1:400 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).