For Research Use Only

PKC Iota Monoclonal antibody

Catalog Number:66493-1-lg Featured Product

2 Publications



Basic Information

Catalog Number: 66493-1-lg Source:

Isotype: IgG2a

Mouse

Immunogen Catalog Number:

AG4990

GenBank Accession Number:

BC022016 GeneID (NCBI): 5584 **UNIPROT ID:** P41743

Full Name: protein kinase C, iota

Calculated MW: 68 kDa Observed MW:

70 kDa

Purification Method:

Protein A purification

CloneNo.: 3A1B6

Recommended Dilutions:

WB: 1:5000-1:50000 IHC: 1:150-1:600 IF-P: 1:200-1:800 IF/ICC: 1:400-1:1600

Applications

Tested Applications:

WB, IHC, IF/ICC, IF-P, ELISA

Cited Applications:

WB, IHC

Species Specificity: human, mouse, rat Cited Species: human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: MCF-7 cells, HeLa cells, HEK-293 cells, ROS1728 cells, C6 cells, LNCaP cells, Jurkat cells, HSC-T6 cells, RAW 264.7 cells, NIH/3T3 cells

IHC: human lung cancer tissue, IF-P: human lung cancer tissue,

IF/ICC: HeLa cells,

Background Information

The atypical protein kinase C isoform PRKC iota (PRKCI) is a member of the protein kinase C (PKC) family of serine/threonine protein kinases. PKC family comprises at least eight members, which are differentially expressed and are involved in a wide variety of cellular processes. PRKC iota is calcium-independent and phospholipiddependent. It is not activated by phorbolesters or diacylglycerol. This kinase can be recruited to vesicle tubular $clusters \, (VTCs) \, by \, direct interaction \, with \, the \, small \, GTP ase \, RAB2, \, where \, this \, kinase \, phosphorylates \, glyceral dehydency is a constant of the small \, GTP and \, constant of the small \, constant \, c$ 3-phosphate dehydrogenase (GAPD/GAPDH) and plays a role in microtubule dynamics in the early secretory pathway. This kinase is found to be necessary for BCL-ABL-mediated resistance to drug-induced apoptosis and therefore protects leukemia cells against drug-induced apoptosis. PRKC iota plays a key role in cell proliferation, differentiation, and carcinogenesis, and it has been shown to be a human oncogene.

Notable Publications

Author	Pubmed ID	Journal	Application
Antja-Voy Hartley	32456215	Int J Mol Sci	WB
Peng Li	40382656	Cell Commun Signal	WB,IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66493-1-lg (PKC lota antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading




WB result of PKC lota antibody (66493-1-lg; 1:30000; incubated at room temperature for 1.5 hours) with sh-Control and sh-PKC lota transfected Hela cells.




Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 66493-1-lg (PKC iota antibody) at dilution of 1:300 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 66493-1-lg (PKC iota antibody) at dilution of 1:300 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HeLa cells using PKC lota antibody (66493-1-lg, Clone: 3A1B6) at dilution of 1:800 and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human lung cancer tissue using PKC lota antibody (66493-1-lg, Clone: 3A1B6) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).