For Research Use Only

NRF2, NFE2L2 Monoclonal antibody

Catalog Number:66504-1-lg Featured Product 121 Publications



Basic Information

Catalog Number: 66504-1-lg Source: Mouse

Isotype: lgG2b

Immunogen Catalog Number:

AG9469

GenBank Accession Number:

BC011558 GeneID (NCBI): 4780 **UNIPROT ID:**

Q16236 Full Name:

nuclear factor (erythroid-derived 2)-

like 2

Calculated MW: 605 aa, 68 kDa Observed MW: 110 kDa

Applications

Tested Applications:

WB, ELISA

Cited Applications: WB, IHC, IF, IP, CoIP, ChIP

Species Specificity: human, mouse Cited Species:

human, mouse, rat, pig, zebrafish

Positive Controls:

WB: MG132 treated HCT 116 cells, MG132 treated NIH/3T3 cells, MG132 treated HepG2 cells

Purification Method:

Protein A purification

Recommended Dilutions:

WB: 1:5000-1:50000

CloneNo.:

1E9E3

Background Information

NRF2, also named as NFE2L2, belongs to the bZIP family and CNC subfamily. It is a transcription activator that binds to antioxidant response (ARE) elements in the promoter regions of target genes. NRF2 is important for the coordinated up-regulation of genes in response to oxidative stress. It may be involved in the transcriptional activation of genes of the beta-globin cluster by mediating enhancer activity of hypersensitive site 2 of the betaglobin locus control region. Nrf2 is a key player in the regulation of genes encoding for many antioxidative response enzymes. The expression of NRF2 may be induced under oxidative stress (PMID:14567983). In lung cancer, Nrf2 activation in malignant cells has been associated with tumor progression and chemotherapy resistance(PMID:20534738). Identifying patients with abnormal NRF2 expression may be important for selection for chemotherapy in NSCLC. As new investigators break into the emerging field of Nrf2 research, confusion regarding the correct migratory pattern of Nrf2 is causing doubts about the accuracy and reproducibility of published results. This letter provides solid evidence that the actually observed molecular weight of Nrf2 is about 70kDa and 95-110 kDa. (PMID: 22703241).

Notable Publications

Author	Pubmed ID	Journal	Application
Jinliang Liu	34630847	Oxid Med Cell Longev	WB
Zi-Chao Wang	36163178	Cell Death Dis	WB
Lei Zhao	34582963	Food Chem Toxicol	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data







MG132 treated NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 66504-1-1g (NRF2, NFE2L2 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.