

For Research Use Only

SMAD3 Monoclonal antibody

Catalog Number: 66516-1-Ig **108 Publications**



Basic Information

Catalog Number:

66516-1-Ig

Concentration:

2000 ug/ml

Source:

Mouse

Isotype:

IgG1

GenBank Accession Number:

NM_005902

GeneID (NCBI):

4088

UNIPROT ID:

P84022

Full Name:

SMAD family member 3

Calculated MW:

48 kDa

Observed MW:

55-60 kDa

Purification Method:

Protein G purification

CloneNo.:

1F10B2

Recommended Dilutions:

WB 1:2000-1:12000

IHC 1:500-1:2000

Applications

Tested Applications:

WB, IHC, ELISA

Cited Applications:

WB, IHC, IF, IP, CoIP, ChIP

Species Specificity:

human, mouse

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HT-1080 cells, NIH/3T3 cells, A431 cells, HEK-293 cells, Jurkat cells, K-562 cells, 4T1 cells

IHC : human stomach cancer tissue, mouse lung tissue

Background Information

SMAD3, also named as hMAD 3 or Mad3, is a 425 amino acid protein, which contains one MH1 domain and one MH2 domain. SMAD3 localizes in the nucleus and cytoplasm. SMAD3 plays an essential role in development and maintenance of self-tolerance and is a critical mediator of the TGF β signaling pathway. SMAD3 is involved in TGF β dependent regulation of steroidogenesis and in T-cell response to TGF β .

Notable Publications

Author	Pubmed ID	Journal	Application
Shaling Li	36169092	Cancer Sci	WB
Bingyu Xie	36179941	Mol Cell Endocrinol	WB
Wen Zhao	34490599	Histol Histopathol	WB,IHC

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66516-1-Ig (SMAD3 antibody) at dilution of 1:6000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.



Immunohistochemical analysis of paraffin-embedded human stomach cancer tissue slide using 66516-1-Ig (SMAD3 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse lung tissue slide using 66516-1-Ig (SMAD3 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).