For Research Use Only

HMGB1 Monoclonal antibody, PBS Only

Catalog Number:66525-1-PBS



Basic Information

Catalog Number:

66525-1-PBS

Size: 1mg/ml Source: Mouse

Isotype: lgG1

Immunogen Catalog Number:

AG1264

Observed MW: 30 kDa

25 kDa

BC003378

GeneID (NCBI):

UNIPROT ID:

Full Name:

P09429

GenBank Accession Number:

high-mobility group box 1 Calculated MW:

Purification Method:

Protein A purification CloneNo.:

1D12A6

Applications

Tested Applications: WB, Indirect ELISA Species Specificity: human, mouse

Background Information

The HMG (high mobility group) proteins are nonhistone chromosomal proteins that is present in almost all eukaryotic cells, and it functions to stabilize NUCLEOSOME formation and acts as a transcription-factor-like protein that regulates the expression of several genes[PMID: 18160415]. Once injury, infection or other inflammatory stimuli, activated macrophages, mature dendritic cells and natural killer cells can secret HMGB1, which act as a crucial cytokine[PMID: 20163887]. HMGB1 also involved in V(D)J recombination by acting as a cofactor of the RAG complex, stimulating cleavage and RAG protein binding at the 23 bp spacer of conserved recombination signal sequences (RSS)[PMID: 19360789]. Act as a Heparin-binding protein that has a role in the extension of neurite-type cytoplasmic processes in developing cells. HMGB1 (high mobility group box 1) modulates gene expression in the nucleus, but certain immune cells secrete HMGB1 as an extracellular Alarmin to signal tissue damage. The nuclear HMGB1 relocalizes to the extracellular milieu in senescent human and mouse cells in culture and in vivo, which $stimulated\ cytokine\ secretion\ through\ TLR-4\ signaling\ (23649808). This\ antibody\ is\ raised\ against\ full\ length\ HMGB1$ of human origin. The monomeric HMGB1 is 29 kDa and dimer is 58 kDa.

Storage


Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS Only

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66525-1-lg (HMGB1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66525-1-PBS in a different storage buffer formulation.