For Research Use Only

SMARCA4/BRG1 Monoclonal antibody

Catalog Number:66561-1-lg Featured Product 2 Publications



Basic Information

Catalog Number: 66561-1-lg Concentration: 1000 ug/ml Source:

Mouse Isotype: lgG2b

Immunogen Catalog Number:

AG16256

GenBank Accession Number:

BC150298 GeneID (NCBI): 6597 **UNIPROT ID:** P51532 Full Name:

SWI/SNF related, matrix associated, actin dependent regulator of chromatin, subfamily a, member 4

Calculated MW: 1647 aa, 185 kDa Observed MW: 185 kDa

Purification Method:

Protein A purification CloneNo.:

2E6B6

Recommended Dilutions: WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total

protein lysate IF/ICC 1:400-1:1600

Applications

Tested Applications: WB, IF/ICC, IP, ELISA **Cited Applications:** WB. IF. ChIP Species Specificity: human, mouse, rat **Cited Species:**

human, mouse

WB: HepG2 cells, COLO 320 cells, HeLa cells, Jurkat cells, K-562 cells, HSC-T6 cells, 4T1 cells

IP: HeLa cells.

IF/ICC: HepG2 cells, PC-3 cells

Background Information

 $SMARCA4, also \ named \ as \ BAF190A, BRG1, SNF2B \ and \ SNF2L4, belongs \ to \ the \ SNF2/RAD54 \ helicase \ family.$ SMARCA4 is a transcriptional coactivator cooperating with nuclear hormone receptors to potentiate transcriptional activation. It is a component of the CREST-BRG1 complex, a multiprotein complex that regulates promoter activation by orchestrating a calcium-dependent release of a repressor complex and a recruitment of an activator $complex. \ It is also involved in vitamin D-coupled transcription regulation via its association with the WINAC involved in vitamin D-coupled transcription regulation via its association with the WINAC involved in vitamin D-coupled transcription regulation via its association with the WINAC involved in vitamin D-coupled transcription regulation via its association with the WINAC involved in vitamin D-coupled transcription regulation via its association with the WINAC involved in vitamin D-coupled transcription regulation via its association with the WINAC involved in vitamin D-coupled transcription regulation via its association with the WINAC involved in vitamin D-coupled transcription regulation via its association with the WINAC involved in vitamin D-coupled transcription regulation via its association with the WINAC involved in vitamin D-coupled transcription regulation via its association via its associati$ complex, a chromatin-remodeling complex recruited by vitamin D receptor (VDR), which is required for the ligandbound VDR-mediated transrepression of the CYP27B1 gene.

Notable Publications

Author	Pubmed ID	Journal	Application
C C Liu	31041569	Inflammation	WB,IF
Tao Ban	39726787	Front Pharmacol	ChIP

Storage


Storage:

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




IP result of anti-SMARCA4/BRG1 (IP:66561-1-Ig, 5ug; Detection:66561-1-Ig 1:3000) with HeLa cells lysate 3200 ug.




WB result of SMARCA4/BRG1 antibody (66561-1-lg; 1:5000; incubated at room temperature for 1.5 hours) with sh-Control and sh-SMARCA4/BRG1 transfected HepG2 cells.




Various lysates were subjected to SDS PAGE followed by western blot with 66561-1-1g (SMARCA4/BRG1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



COLO 320 cells were subjected to SDS PAGE followed by western blot with 66561-1-1g (SMARCA4/BRG1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using SMARCA4/BRG1 antibody (66561-1-lg, Clone: 2E6B6) at dilution of 1:50000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed PC-3 cells using SMARCA4/BRG1 antibody (66561-1-lg, Clone: 2E6B6) at dilution of 1:50000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using SMARCA4/BRG1 antibody (66561-1-Ig, Clone: 2E6B6) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L), CL594-Phalloidin (red).