For Research Use Only

NMDAR2B/GRIN2B Monoclonal antibody

Proteintech®
Antibodies | ELISA kits | Proteins
www.ptglab.com

Purification Method:

CloneNo.:

1C5E12

Protein G purification

Recommended Dilutions:

WB 1:1000-1:6000

Catalog Number:66565-1-lg

1 Publications

Basic Information

Applications

Catalog Number: 66565-1-Ig Size: 1300 µg/ml

Source: Mouse Isotype: IgG1

Immunogen Catalog Number:

AG16718

Tested Applications:

WB, ELISA

Cited Applications:

WB

Species Specificity: Human, Mouse, Rat Cited Species: mouse GenBank Accession Number:

BC113620 GeneID (NCBI): 2904 UNIPROT ID:

Q13224 Full Name:

glutamate receptor, ionotropic, N-methyl D-aspartate 2B

Calculated MW: 1484 aa, 166 kDa

Observed MW: 166 kDa

Positive Controls:

WB: mouse brain tissue, rat brain tissue

Background Information

GRIN2B (also known as GluN2B or NMDAR2B) is a member of the N-methyl-D-aspartate (NMDA) receptor family within the ionotropic glutamate receptor superfamily. NMDA receptors are widely expressed in the central nervous system and play a major role in excitatory synaptic transmission and plasticity (PMID: 23223336). NMDA receptors large multi-subunit complexes arranged into heteromeric assemblies composed of four homologous subunits within a repertoire of over 10 different subunits: eight GluN1 isoforms, four GluN2 subunits (A-D) and two GluN3 subunits (A and B) (PMID: 21395862). Naturally occurring mutations within GRIN2B gene are associated with neurodevelopmental disorders including autism spectrum disorder, attention deficit hyperactivity disorder, epilepsy, and schizophrenia.

Notable Publications

Author	Pubmed ID	Journal	Application
Rasa Volianskis	38853552	Philos Trans R Soc Lond B Biol Sci	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

Selected Validation Data



mouse brain tissue were subjected to SDS PAGE followed by western blot with 66565-1-lg (NMDAR2B/GRIN2B antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.