For Research Use Only

CA3 Monoclonal antibody

Catalog Number:66608-1-Ig



Basic Information

Catalog Number: 66608-1-lg

 Size:
 GeneID (NCBI):

 1600 μg/ml
 761

 Source:
 UNIPROT ID:

Source: UNIPROT ID:

Mouse P07451

Isotype: Full Name:
IgG2b carbonic anhydrase III, muscle

Immunogen Catalog Number:

AG7513

29 kDa Observed MW: 29 kDa Purification Method: Protein A purification

CloneNo.: 3C10A2

Recommended Dilutions: WB 1:20000-1:100000 IHC 1:500-1:2000 IF 1:200-1:800

Applications

Tested Applications: WB, IF-P, IHC, ELISA Species Specificity: Human, Mouse, Rat, Pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0 Positive Controls:

WB: pig skeletal muscle tissue, rat skeletal muscle tissue, mouse skeletal muscle tissue

IHC: mouse skeletal muscle tissue, IF: mouse skeletal muscle tissue,

Background Information

Carbonic anhydrase III (CA3), which belongs to the alpha-carbonic anhydrase family, is a cytoplasmic enzyme that exhibits a relatively low carbon dioxide hydratase activity. It is expressed at a very high level in skeletal muscle, where physical exercise has been shown to increase free radical production. In addition to its carbon dioxide hydratase activity, CA3 has been demonstrated to have a carboxyl esterase activity and phosphatase activity, which suggests that it is a tyrosine phosphatase (PMID: 10064618). CA3 was found to be localized in Type-I muscle fibers and could be used as a marker for abnormal Type-I muscle fibers in several neuromuscular diseases (PMID: 6221502).

GenBank Accession Number:

BC004897

specific

Calculated MW:

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66608-1-lg (CA3 antibody) at dilution of 1:100000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse skeletal muscle tissue slide using 66608-1-Ig (CA3 antibody) at dilution of 1:1000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse skeletal muscle tissue slide using 66608-1-Ig (CA3 antibody) at dilution of 1:1000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded mouse skeletal muscle tissue using CA3 antibody (66608-1-lg, Clone: 3C10A2) at dilution of 1:400 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).