For Research Use Only

CCRL2 Monoclonal antibody

Catalog Number:66611-1-lg 4 Publications



Basic Information

Catalog Number: 66611-1-lg

BC025717 GeneID (NCBI): 9034

GenBank Accession Number:

Purification Method: Protein G purification

CloneNo.: Size: 1300 µg/ml 2C1E11 **UNIPROT ID:** Recommended Dilutions: Source:

Mouse 000421 WB 1:1000-1:6000 IHC 1:400-1:1600 Full Name: Isotype:

lgG1 chemokine (C-C motif) receptor-like 2

Calculated MW: Immunogen Catalog Number: AG4045 344 aa, 40 kDa Observed MW:

47 kDa

Applications

Tested Applications: IHC, WB,ELISA **Cited Applications:**

WB

Species Specificity:

Human **Cited Species:** human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, K-562 cells IHC: human spleen tissue,

Background Information

Chemokine CC motif receptor-like 2 (CCRL2) is a seven-transmembrane domain receptor that shares structural and functional similarities with the family of atypical chemokine receptors (ACKRs) (PMID: 28743719). CCRL2 does not appear to be a signaling receptor, but may have a role in modulating chemokine-triggered immune responses by capturing and internalizing CCL19 or by presenting RARRES2 ligand to CMKLR1, a functional signaling receptors. CCRL2 has been found on almost all hemopoietic cells, including monocytes, macrophages, polymorphonuclear neutrophils (PMNs), T cells, dendritic cells, natural killer cells, and CD34+ progenitor cells, with PMNs expressing the highest levels (PMID: 15188357). CCRL2 is also expressed by barrier cells, such as lymphatic and blood endothelial cells and bronchial and intestinal epithelial cells (PMID: 29056935).

Notable Publications

Author	Pubmed ID	Journal	Application
Deyu Li	31853188	Onco Targets Ther	WB
Ling Ding	31830143	PLoS Pathog	WB
Lunbo Tan	38361240	Hypertension	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



HeLa cells and A431 cells were subjected to SDS PAGE followed by western blot with 66611-1-1g (CCRL2 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human spleen tissue slide using 66611-1-lg (CCRL2 antibody) at dilution of 1:800 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human spleen tissue slide using 66611-1-Ig (CCRL2 antibody) at dilution of 1:800 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).