For Research Use Only

RABEPK/p40 Monoclonal antibody

Catalog Number:66622-1-lg 1 Publications



Basic Information

Catalog Number:

GenBank Accession Number: BC065725

Purification Method:

66622-1-lg Size:

GeneID (NCBI):

Protein A purification

1000 µg/ml

10244

CloneNo.: 1E11A3

Source:

Full Name: Rab9 effector protein with kelch Recommended Dilutions:

Mouse Isotype:

motifs

WB 1:1000-1:6000 IHC 1:500-1:2000

lgG1

Calculated MW:

41 kDa

Immunogen Catalog Number:

IF 1:50-1:500

AG7796

Observed MW: 40 kDa

Applications

Tested Applications:

IF, IHC, WB, ELISA

Cited Applications:

Positive Controls:

WB: Jurkat cells, HeLa cells, HEK-293 cells, HSCT6

cells,RAW 264.7 cells

IHC: human lung cancer tissue, human colon tissue

IF: A549 cells,

Species Specificity: Human, mouse, rat **Cited Species:**

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

Rab9 GTPase is required for the transport of mannose 6-phosphate receptors from endosomes to the trans-Golgi network in living cells, and in an in vitro system that reconstitutes this process. P40 is an effector of Rab9 that interacts preferentially with the active form of Rab9. p40 does not interact with Rab7 or K-Ras; it also fails to bind Rab9 when it is bound to GDI. The protein is found in cytosol, yet a significant fraction (~30%) is associated with $cellular\ membranes.\ P40\ is\ a\ very\ potent\ transport\ factor\ in\ that\ the\ pure,\ recombinant\ protein\ can\ stimulate,$ significantly, an in vitro transport assay that measures transport of mannose 6-phosphate receptors from endosomes to the trans-Golgi network.

Notable Publications

Author	Pubmed ID	Journal	Application
Jun Fu	35949347	Exp Ther Med	IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Jurkat cells were subjected to SDS PAGE followed by western blot with 66622-1-1g (RABEPK/p40 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 66622-1-Ig (RABEPK/p40 antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 66622-1-Ig (RABEPK/p40 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed A549 cells using 66622-1-lg (RABEPK/p40 antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).