For Research Use Only

p70(S6K) Monoclonal antibody


Catalog Number:66638-1-lg 13 Publications



Basic Information	Catalog Number: 66638-1-Ig	GenBank Accession Number: BC053365	Purification Method: Protein G purification	
	Concentration:	GenelD (NCBI):	CloneNo.:	
	1200 µg/ml	6198	1D11C6	
	Source: Mouse	UNIPROT ID: P23443	Recommended Dilutions: WB 1:1000-1:6000	
	Isotype:	Full Name:	IHC 1:200-1:800	
	lgG1 ribosomal protein S6 kinase, 70kDa,		Da,	
	Immunogen Catalog Number:	polypeptide 1 Calculated MW: 59 kDa		
	AG5883			
		Observed MW: 60-75 kDa		
Applications	Tested Applications: WB, IHC, ELISA	Positive Controls:		
	Cited Applications:	WB : Jurka cells		
	WB, IF			
	Species Specificity: Human, Mouse, Rat		in lung cancer tissue,	
	Cited Species: human, mouse			
	Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0			
	RPS6KB1(Ribosomal protein S6 kinase beta-1) is also named as STK14A, p70 S6KA and belongs to the S6 kinase subfamily. RPS6KB1 is a major substrate of MTOR and acts as a crucial effector of MTOR signaling pathway. It plays a key role in cell growth and proliferation by regulating INS sensitivity, metabolism, protein synthesis, and cell cycle. RPS6KB1 may play an important role in the progression of HCC and could serve as a potential molecular target for HCC therapy (PMID:22684641). RPS6KB1 is a 70 kDa protein and has 5 isoforms with the calculated molecular mass of 51-59kDa produced by alternative initiation.			
Background Information	subfamily. RPS6KB1 is a major sub key role in cell growth and prolife RPS6KB1 may play an important ro HCC therapy (PMID:22684641). RP	ration by regulating INS sensitivity, n ole in the progression of HCC and coul 'S6KB1 is a 70 kDa protein and has 5 is	ffector of MTOR signaling pathway. It plays netabolism, protein synthesis, and cell cycle d serve as a potential molecular target for	
	subfamily. RPS6KB1 is a major sub key role in cell growth and prolife RPS6KB1 may play an important ro HCC therapy (PMID:22684641). RP of 51-59kDa produced by alternati	ration by regulating INS sensitivity, n ole in the progression of HCC and coul 'S6KB1 is a 70 kDa protein and has 5 is	Frector of MTOR signaling pathway. It plays letabolism, protein synthesis, and cell cycle d serve as a potential molecular target for oforms with the calculated molecular mass	
	subfamily. RPS6KB1 is a major sub key role in cell growth and prolife RPS6KB1 may play an important rr HCC therapy (PMID:22684641). RP of 51-59kDa produced by alternati	ration by regulating INS sensitivity, n ole in the progression of HCC and coul 'S6KB1 is a 70 kDa protein and has 5 is ive initiation.	ffector of MTOR signaling pathway. It plays a netabolism, protein synthesis, and cell cycle d serve as a potential molecular target for	
Background Information Notable Publications	subfamily. RPS6KB1 is a major sub key role in cell growth and prolife RPS6KB1 may play an important ro HCC therapy (PMID:22684641). RP of 51-59kDa produced by alternati Author Haitao Fan	ration by regulating INS sensitivity, n ole in the progression of HCC and coul IS6KB1 is a 70 kDa protein and has 5 is ive initiation. Pubmed ID Journal	Frector of MTOR signaling pathway. It plays netabolism, protein synthesis, and cell cycle d serve as a potential molecular target for oforms with the calculated molecular mass Application WB	


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





Various lysates were subjected to SDS PAGE followed by western blot with 66638-1-lg (p70(S6K) antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 66638-1-Ig (p70(S6K) antibody) at dilution of 1:400 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human lung cancer tissue slide using 66638-1-Ig (p70(S6K) antibody) at dilution of 1:400 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).