For Research Use Only

TLR2 Monoclonal antibody, PBS Only

Catalog Number: 66645-1-PBS

Featured Product



Purification Method:

Protein A purification

CloneNo.:

1C1D3

Basic Information

Catalog Number: 66645-1-PBS

Size: 1 mg/ml GenBank Accession Number:

BC033756

GeneID (NCBI):

7097

Source: UNIPROT ID: Mouse 060603

Isotype: Full Name:
IgG2a toll-like receptor 2

Immunogen Catalog Number: Calculated MW:

AG25626 784 aa, 90 kDa

Observed MW:

90-100 kDa

Applications

Tested Applications:

WB, IF, IHC, Indirect ELISA

Species Specificity:

Human


Background Information

TLR2, also known as CD282, is a member of the Toll-like receptor (TLR) family. TLR receptors are type I integral transmembrane glycoproteins which play a fundamental role in pathogen recognition and activation of innate immunity. TLR2 is expressed by immune cells including monocytes/macrophages, polymorphonuclear leukocytes (PMLs) and dendritic cells, and is also expressed by some no-immune cells including endothelial and epithelial cells (PMID: 21428911; 22566960). TLR2 forms heterodimers with TLR1 and TLR6, which is the initial step in a cascade of events leading to significant innate immune responses, development of adaptive immunity to pathogens and protection from immune sequelae related to infection with these pathogens (PMID: 22566960).


Storage

Storage: Store at -80°C. Storage Buffer: PBS Only


Selected Validation Data




WB result of TLR2 antibody (66645-1-lg; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-TLR2 transfected THP-1 cells. This data was developed using the same antibody clone with 66645-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human liver tissue slide using 66645-1-Ig (TLR2 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66645-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed THP-1 cells using TLR2 antibody (66645-1-1g, Clone: 1C1D3) at dilution of 1:400 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66645-1-PBS in a different storage buffer formulation



THP-1 cells were subjected to SDS PAGE followed by western blot with 66645-1-1g (TLR2 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66645-1-PBS in a different storage buffer formulation.