### For Research Use Only

# PARK2/Parkin Monoclonal antibody

Catalog Number:66674-1-lg Featured Product

64 Publications



**Basic Information** 

Catalog Number: 66674-1-lg Source: Mouse

Isotype: lgG2b

Immunogen Catalog Number:

AG5179

GenBank Accession Number:

BC022014 GeneID (NCBI): 5071 **UNIPROT ID:** 060260

Parkinson disease (autosomal recessive, juvenile) 2, parkin

Calculated MW: 52 kDa Observed MW: 42-52 kDa

Full Name:

**Purification Method:** 

Protein A purification

CloneNo.: 2H5A7

Recommended Dilutions: WB: 1:2000-1:8000 IHC: 1:250-1:1000

IF/ICC: 1:400-1:1600

**Applications** 

**Tested Applications:** WB, IHC, IF/ICC, ELISA **Cited Applications:** WB, IHC, IF

Species Specificity: human, mouse Cited Species:

human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HEK-293 cells, SH-SY5Y cells, U-251 cells, mouse

brain tissue

IHC: mouse brain tissue, IF/ICC: RAW 264.7 cells,

# **Background Information**

Parkin, a RING-type E3 ubiquitin-protein ligase, is involved in the ubiquitination pathway and contributes to protection from neurotoxicity induced by unfolded protein stresses. Its ubiquitin-protein ligase activity promotes the degradation of a viariety of proteins including itself. Mutations in Parkin are implicated in the pathogenesis of autosomal recessive familial Parkinson's disease. It has 8 isoforms produced by alternative splicing.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Tetsushi Hirano	34520793	Toxicol Appl Pharmacol	WB
Xue Wang	32963699	Oxid Med Cell Longev	IF
Huijuan Xiao	36308405	J Cell Mol Med	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage


## **Selected Validation Data**




HEK-293 cells were subjected to SDS PAGE followed by western blot with 66674-1-1g (Parkin antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 66674-1-Ig (Parkin antibody) at dilution of 1:500 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 66674-1-Ig (Parkin antibody) at dilution of 1:500 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed RAW 264.7 cells using PARK2/Parkin antibody (66674-1-lg, Clone: 2H5A7) at dilution of 1:400 and CoraLite® 488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L) (SA00013-1), CL594-Phalloidin (red).