For Research Use Only

HMG20A Monoclonal antibody, PBS Only




Catalog Number:66676-1-PBS


Basic Information	Catalog Number: 66676-1-PBS	GenBank Accession Number: BC021959	Purification Method: Protein G purification
	Concentration: 1mg/ml	GenelD (NCBI): 10363	CloneNo.: 1B10F12
	Source: Mouse	UNIPROT ID: Q9NP66	
	lsotype: lgG1	Full Name: high-mobility group 20A	
	Immunogen Catalog Number: AG2718	Calculated MW: 347 aa, 40 kDa	
		Observed MW: 45-50 kDa	
Applications	Tested Applications: WB, IF/ICC, Indirect ELISA		
	Species Specificity: human, mouse		
Background Information	The HMG (high mobility group) proteins are nonhistone chromosomal proteins that is present in almost all eukaryotic cells, and it functions to stabilize nucleosome formation and acts as a transcription-factor-like protein that regulates the expression of several genes. HMG20A has a role in the initiation of neuronal differentiation by activating REST-responsive genes, and it also nvolves in the recruitment of the histone methyltransferase MLL and consequent increased methylation of histone H3 lysine 4. HMG20A (for inhibitor of BRAF35) overcomes the repressive effects of the neuronal silencer REST and activates REST-responsive genes through the modulation of histone methylation		
Storage	Storage: Store at -80°C. The product is shipped with ice pa Storage Buffer:	cks. Upon receipt, store it immediatel	.y at -80°C

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





HeLa cells were subjected to SDS PAGE followed by western blot with 66676-1-1g (HMG20A antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66676-1-PBS in a different storage buffer formulation.

Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 66676-1-1g (HMG20A antibody) at dilution of 1:100 and Alexa Fluor 488-conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66676-1-PBS in a different storage buffer formulation.