

For Research Use Only

# B3GAT1 Monoclonal antibody, PBS Only

Catalog Number: 66711-1-PBS



## Basic Information

Catalog Number:

66711-1-PBS

Size:

1 mg/ml

Source:

Mouse

Isotype:

IgG2a

Immunogen Catalog Number:

AG13593

GenBank Accession Number:

BC010466

GeneID (NCBI):

27087

UNIPROT ID:

Q9P2W7

Full Name:

beta-1,3-glucuronyltransferase 1  
(glucuronosyltransferase P)

Calculated MW:

38 kDa

Observed MW:

38 kDa, 100 kDa

Purification Method:

Protein A purification

CloneNo.:

1F4A11

## Applications

Tested Applications:

WB, FC (Intra), ELISA

Species Specificity:

human, mouse

## Background Information

B3GAT1 is associated with glycosaminoglycan biosynthesis. It plays a key role in neurodevelopment. The calculated molecular weight of B3GAT1 is 38 kDa and 66711-1-Ig antibody detects 100 kDa form in SDS-PAGE which might as a result of homodimer or glycosylation.

## Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66711-1-Ig (B3GAT1 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66711-1-PBS in a different storage buffer formulation.



mouse brain tissue were subjected to SDS PAGE followed by western blot with 66711-1-Ig (B3GAT1 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66711-1-PBS in a different storage buffer formulation.



1X10<sup>6</sup> K-562 cells were intracellularly stained with 0.4 ug Anti-Human CD57 (66711-1-Ig, Clone:1F4A11) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2a Isotype Control (66360-2-Ig, Clone: K11A1B2A2) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 66711-1-PBS in a different storage buffer