

For Research Use Only

# XIAP Monoclonal antibody

Catalog Number: 66800-1-Ig

Featured Product

12 Publications



## Basic Information

Catalog Number:

66800-1-Ig

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG18088

GenBank Accession Number:

BC032729

GeneID (NCBI):

331

UNIPROT ID:

P98170

Full Name:

X-linked inhibitor of apoptosis

Calculated MW:

60 kDa

Observed MW:

57 kDa

Purification Method:

Protein G purification

CloneNo.:

2A1A12

Recommended Dilutions:

WB: 1:5000-1:50000

IHC: 1:250-1:1000

IF-P: 1:200-1:800

FC (Intra): 0.40 ug per 10<sup>6</sup> cells in a 100 µl suspension

## Applications

Tested Applications:

WB, IHC, IF-P, FC (Intra), ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

human, mouse, rat

Cited Species:

human, mouse

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

Positive Controls:

WB: LNCaP cells, HEK-293 cells, HeLa cells, HepG2 cells, HEK-293T cells, Jurkat cells

IHC: human colon tissue, human lung cancer tissue

IF-P: human colon cancer tissue, human colon tissue

FC (Intra): HeLa cells,

## Background Information

XIAP, also named as API3, BIRC4 and IAP3, belongs to the IAP family. It has E3 ubiquitin-protein ligase activity. It mediates the proteasomal degradation of target proteins, such as caspase-3, SMAC or AIFM1. XIAP is an inhibitor of caspase-3, -7 and -9. It mediates activation of MAP3K7/TAK1, leading to the activation of NF-kappa-B. XIAP is an apoptotic suppressor. It is ubiquitinated and degraded by the proteasome in apoptotic cells.

## Notable Publications

Author	Pubmed ID	Journal	Application
Jingwen Tan	36208777	Chem Biol Interact	WB
Siwen Yin	35154432	Oncol Lett	WB
Yuhui Liu	33613115	Int J Biol Sci	WB

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66800-1-Ig (XIAP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control.




WB result of XIAP antibody (66800-1-Ig; 1:5000; incubated at room temperature for 1.5 hours) with sh-Control and sh-XIAP transfected HEK-293 cells.




Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 66800-1-Ig (XIAP antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 66800-1-Ig (XIAP antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human colon cancer tissue using XIAP antibody (66800-1-Ig, Clone: 2A1A12) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



1X10<sup>6</sup> HeLa cells were intracellularly stained with 0.4 ug Anti-Human XIAP (66800-1-Ig, Clone:2A1A12) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer.