

For Research Use Only

XIAP Monoclonal antibody, PBS Only

Catalog Number: 66800-1-PBS

Featured Product



Basic Information

Catalog Number:

66800-1-PBS

Size:

1mg/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG18088

GenBank Accession Number:

BC032729

GeneID (NCBI):

331

UNIPROT ID:

P98170

Full Name:

X-linked inhibitor of apoptosis

Calculated MW:

60 kDa

Observed MW:

57 kDa

Purification Method:

Protein G purification

CloneNo.:

2A1A12

Applications

Tested Applications:

WB, IF, IHC, Indirect ELISA

Species Specificity:

Human, mouse, rat

Background Information

XIAP, also named as API3, BIRC4 and IAP3, belongs to the IAP family. It has E3 ubiquitin-protein ligase activity. It mediates the proteasomal degradation of target proteins, such as caspase-3, SMAC or AIFM1. XIAP is an inhibitor of caspase-3, -7 and -9. It mediates activation of MAP3K7/TAK1, leading to the activation of NF-kappa-B. XIAP is an apoptotic suppressor. It is ubiquitinated and degraded by the proteasome in apoptotic cells.

Storage

Storage:

Store at -80°C.

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer:

PBS Only

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.



Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66800-1-Ig (XIAP antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated GAPDH Monoclonal antibody (HRP-60004) as loading control. This data was developed using the same antibody clone with 66800-1-PBS in a different storage buffer formulation.




WB result of XIAP antibody (66800-1-Ig; 1:5000; incubated at room temperature for 1.5 hours) with sh-Control and sh-XIAP transfected HEK-293 cells. This data was developed using the same antibody clone with 66800-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 66800-1-Ig (XIAP antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66800-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed human colon cancer tissue using XIAP antibody (66800-1-Ig, Clone: 2A1A12) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 66800-1-PBS in a different storage buffer formulation.



1X10⁶ HeLa cells were intracellularly stained with 0.4 ug Anti-Human XIAP (66800-1-Ig, Clone:2A1A12) and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C). This data was developed using the same antibody clone with 66800-1-PBS in a different storage buffer formulation.