For Research Use Only

CCR6 Monoclonal antibody

Catalog Number:66801-1-lg 2 Publications



Basic Information

Catalog Number: 66801-1-lg

Size:

BC037960 GeneID (NCBI): 1235 **UNIPROT ID:**

P51684

Full Name:

GenBank Accession Number:

chemokine (C-C motif) receptor 6

Purification Method: Protein A purification CloneNo.:

1C11A9

1600 µg/ml Source: Mouse Isotype: IgG2a

Recommended Dilutions: IHC 1:250-1:1000 IF-P 1:400-1:1600 IF/ICC 1:400-1:1600

Immunogen Catalog Number:

AG24195 42 kDa

Calculated MW:

Applications

Tested Applications: IF/ICC,IF-P, IHC, ELISA Cited Applications:

IF, IHC

Species Specificity:

Human **Cited Species:** human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

IHC: human tonsillitis tissue, IF-P: human tonsillitis tissue,

IF/ICC: Jurkat cells, human tonsillitis tissue

Background Information

CCR6, also known as CD196, is a C-C chemokine receptor that belongs to family A of G protein-coupled receptor superfamily. It is expressed in most B cells, memory T cells, and some subsets of dendritic cells (PMID: 17057190). CCR6 is a receptor for the C-C type chemokine CCL20 (PMID: 9169459). The ligand-receptor pair CCL20-CCR6 is $responsible for the chemoattraction of immature dendritic cells, effector/memory \ T-cells, and \ B-cells \ and \ plays \ and \ an$ role in skin and mucosal surfaces under homeostatic and inflammatory conditions, as well as in pathology, including cancer and rheumatoid arthritis (PMID: 12948524). CCR6 polymorphism has been associated with rheumatoid arthritis susceptibility and CCR6 is involved in IL-17-driven autoimmunity in human diseases (PMID: 20453841).

Notable Publications

Author	Pubmed ID	Journal	Application
Jing Ouyang	38902257	Cell Death Dis	IF
Yufei Liu	37579688	Neoplasia	IHC

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Immunofluorescent analysis of (4% PFA) fixed Jurkat cells using CCR6 antibody (66801-1-lg, Clone: 1C11A9) at dilution of 1:800 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66801-1-1g (CCR6 antibody) at dilution of 1:500 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 66801-1-1g (CCR6 antibody) at dilution of 1:500 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CCR6 antibody (66801-1-lg, Clone: 1C11A9) at dilution of 1:800 and Coralite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).