

For Research Use Only

OGT Monoclonal antibody

Catalog Number: 66823-1-Ig

Featured Product

7 Publications



Basic Information

Catalog Number:

66823-1-Ig

Size:

1400 µg/ml

Source:

Mouse

Isotype:

IgG2a

Immunogen Catalog Number:

AG28402

GenBank Accession Number:

BC014434

GeneID (NCBI):

8473

UNIPROT ID:

O15294

Full Name:

O-linked N-acetylglucosamine (GlcNAc) transferase (UDP-N-acetylglucosamine:polypeptide-N-acetylglucosaminyl transferase)

Calculated MW:

1046 aa, 117 kDa

Observed MW:

110 kDa

Purification Method:

Protein A purification

CloneNo.:

2B2A6

Recommended Dilutions:

WB 1:5000-1:50000

IHC 1:250-1:1000

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, ELISA

Cited Applications:

WB, IHC, IF

Species Specificity:

Human, mouse, rat, pig

Cited Species:

human, mouse

Positive Controls:

WB: HeLa cells, HEK-293 cells, HSC-T6 cells, NIH/3T3 cells, pig brain tissue, rat brain tissue, mouse brain tissue

IHC: human lung cancer tissue, human colon cancer tissue

IF/ICC: HepG2 cells,

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Background Information

O-linked N-acetylglucosamine transferase (OGT) catalyzes the attachment of N-acetylglucosamine (GlcNAc) monosaccharides to the hydroxyl group of serine or threonine residues of numerous nuclear and cytoplasmic proteins and may play important roles in a large number of diverse intracellular processes ranging from translational control, transcription, transcriptional repression, INS resistance and regulation of the cell cycle. It exists as a heterotrimeric complex with two 110 kDa and one 70 kDa subunits. Recent studies have shown that O-GlcNAcylation plays essential roles in cancer formation and progression. O-GlcNAcylation as well as OGT expression was found to be significantly elevated in the cancer tissues.

Notable Publications

Author	Pubmed ID	Journal	Application
Bo Xu	35690146	J Biol Chem	WB
S I Panpan	39496915	Mol Cell Biochem	WB, IHC
Stephen Henry Holland	39456185	Biomolecules	WB

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com




W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 66823-1-Ig (OGT antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using OGT antibody (66823-1-Ig, Clone: 2B2A6) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L), CL594-phalloidin (red).