## For Research Use Only

## NeuN Monoclonal antibody, PBS Only

Catalog Number:66836-1-PBS



**Basic Information** 

Catalog Number:

GenBank Accession Number: NM\_001082575

Purification Method:

66836-1-PBS

GeneID (NCBI):

Protein A purification

Size: 1mg/ml

146713

CloneNo.: 3A4C1

Source:

Full Name: hexaribonucleotide binding protein 3

Mouse Isotype:

lgG1

Immunogen Catalog Number:

AG28016

**Applications** 

Tested Applications: IHC, IF-P, FC (Intra), ELISA Species Specificity:

human, mouse, rat

## **Background Information**

NeuN, encoded by FOX3, is a neuron-specific nuclear protein. Anti-NeuN stains exclusively neuronal cells in the central and peripheral nervous systems, especially postmitotic and differentiating neurons, as well as terminally differentiated neurons. Anti-NeuN has been used widely as a reliable tool to detect most postmitotic neuronal cell types. The immunohistochemical staining is primarily localized in the nucleus of the neurons with lighter staining in the cytoplasm.

Storage

Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C

Storage Buffer: PBS Only


## Selected Validation Data




Immunohistochemical analysis of paraffinembedded rat cerebellum tissue slide using 66836-1-Ig (NeuN antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Sodium Citrate buffer (pH 6.0). This data was developed using the same antibody clone with 66836-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (4% PFA) fixed rat cerebellum tissue using 66836-1-lg (NeuN antibody, red), at dilution of 1:200 and CoraLite®594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). The section was co-stained with 14479-1-AP (Calbindin-D28k antibody, green). This data was developed using the same antibody clone with 66836-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded rat cerebellum tissue slide using 66836-1-Ig (NeuN antibody) at dilution of 1:5000 (under 40x lens). Heat mediated antigen retrieval with Sodium Citrate buffer (pH 6.0). This data was developed using the same antibody clone with 66836-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (4% PFA) fixed mouse cerebellum tissue using 66836-1-Ig (NeuN antibody), at dilution of 1:100 and Coralite®594-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). The section was co-stained with 14479-1-AP (Calbindin-D28k Antibody, green). This data was developed using the same antibody clone with 66836-1-PBS in a different storage buffer formulation.



1X10^6 SH-SY5Y cells were intracellularly stained with 0.2 ug Anti-Human NeuN (66836-1-lg, Clone:3A4C1) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgC(H+L) at dilution 1:1000 (red), or 0.2 ug Control Antibody. Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011). This data was developed using the same antibody clone with 66836-1-PBS in a different storage buffer formulation.



Immunohistochemical analysis of paraffinembedded human brain tissue slide using 66836-1-Ig (NeuN antibody) at dilution of 1:20000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66836-1-PBS in a different storage buffer formulation.