For Research Use Only

CPLX2 Monoclonal antibody

Catalog Number: 66849-1-Ig



Basic Information

Catalog Number: GenBank Accession Number: 66849-1-lg BC093706

 66849-1-Ig
 BC093706

 Size:
 GeneID (NCBI):

 1900 ug/ml
 10814

 Source:
 UNIPROT ID:

MouseQ6PUV4Isotype:Full Name:IgG1complexin 2Immunogen Catalog Number:Calculated MW:AG27949134 aa, 15 kDa

Observed MW: 18-20 kDa

Applications

Tested Applications: WB, IHC, FC (Intra), ELISA Species Specificity:

human, mouse, rat, pig

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate

buffer pH 6.0

Purification Method: Protein G purification

CloneNo.: 1B2A11

Positive Controls:

brain tissue, rat cerebellum tissue

IHC: mouse brain tissue,

Recommended Dilutions: WB 1:500-1:3000 IHC 1:250-1:1000

WB: mouse brain tissue, mouse cerebellum tissue, rat

Background Information

Complexins are soluble proteins that regulate the activity of soluble N-ethylmaleimide-sensitive factor attachment protein receptor (SNARE) complexes necessary for vesicle fusion. Neuronal specific complexin 1 (CPLX1) has inhibitory and stimulatory effects on exocytosis by clamping trans-SNARE complexes in a prefusion state and promoting conformational changes to facilitate membrane fusion following cell stimulation. Complexin2 (CPLX2) is a pre-synaptic protein believed to regulate neurotransmitter release from pre-synaptic terminals, it is downregulated in schizophrenic patients suffering from depression, animal models of depression and neurological disorders such as Huntington's disease in which depression is a major symptom.

Storage


Storage:

Store at -20°C. Stable for one year after shipment. Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




mouse brain tissue were subjected to SDS PAGE followed by western blot with 66849-1-1g (CPLX2 antibody) at dilution of 1:1500 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 66849-1-Ig (CPLX2 antibody) at dilution of 1:500 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 66849-1-Ig (CPLX2 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1x10^6 SH-SY5Y cells were intracellularly stained with 0.4 ug CPLY2 Monoclonal antibody (66849-1-1g, Clone:1B2A11) and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(IH-L) (SA00013-1) (red), or 0.4 ug Mouse IgG1 isotype control Mouse McAb (66360-1-1g, Clone: 1F8D3) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).