For Research Use Only

SREBF1 Monoclonal antibody, PBS Only

Catalog Number:66875-1-PBS Featured Product



Basic Information

Catalog Number:

GenBank Accession Number: BC063281

Purification Method: Protein G purification

66875-1-PBS

GeneID (NCBI):

Size: 1mg/ml

6720 **UNIPROT ID:** CloneNo.: 1B6G5

Source: Mouse

P36956 Full Name:

Isotype: lgG1

sterol regulatory element binding

transcription factor 1

AG5484

Calculated MW: 1177 aa. 125 kDa

Observed MW: 125 kDa

Applications

Tested Applications:

Immunogen Catalog Number:

WB, IHC, IF/ICC, ELISA

Species Specificity:

Background Information

SREBF1, also named as BHLHD1 and SREBP1, contains one basic helix-loop-helix (bHLH) domain and belongs to the SREBP family. It is a transcriptional activator required for lipid homeostasis. The SREBPs are synthesized as precursors anchored to endoplasmic reticulum (ER) membranes and complexed with SCAP. When the cellular cholesterol level is low, SREBP-SCAP complexes move to the Golgi apparatus, where SREBPs undergo a two-step proteolytic processing, leading to the release of the mature form, an N-terminal fragment, i.e, basic helix-loop-helix leucine zipper transcription factor. These factors enter the nucleus where they bind to sterol regulatory elements (SRE) in the promoter regions of a number of genes whose products mediate the synthesis of cholesterol and fatty acids. [PMID: 21698267]. This antibody can recognize the 125kd precursor form and the 68kd mature form of human

Storage


Storage:

The product is shipped with ice packs. Upon receipt, store it immediately at -80°C


Storage Buffer:

PBS Only


Selected Validation Data




Immunohistochemical analysis of paraffinembedded human kidney tissue slide using 66875-1-Ig (SREBF1 antibody) at dilution of 1:200 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66875-1-PBS in a different storage buffer formulation.




Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using SREBF1 antibody (66875-1-Ig, Clone: 1B6G5) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). CL594-Phalloidin stains the F-actin (red). DAPI (blue). This data was developed using the same antibody clone with 66875-1-PBS in a different storage buffer formulation.




WB result of SREBF 1 antibody (66875-1-Ig; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-SREBF 1 transfected HeLa cells. This data was developed using the same antibody clone with 66875-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 66875-1-lg (SREBF1 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66875-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 66875-1-1g (SREBF1 antibody) at dilution of 1:8000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66875-1-PBS in a different storage buffer formulation.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using SREBF1 antibody (66875-1-Ig, Clone: 1B6G5) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM002). This data was developed using the same antibody clone with 66875-1-PBS in a different storage buffer formulation.