For Research Use Only

S6 Ribosomal protein Monoclonal antibody



Catalog Number:66886-1-lg

10 Publications

Basic Information

Catalog Number: 66886-1-lg Size:

1500 µg/ml Source: Mouse Isotype: IgG2a

Immunogen Catalog Number:

AG6599

Calculated MW: 29 kDa Observed MW:

29-32 kDa

BC000524

6194

P62753

GeneID (NCBI):

UNIPROT ID:

Full Name:

ribosomal protein S6

GenBank Accession Number:

Purification Method:

Protein A purification CloneNo.:

1C3E10 Recommended Dilutions: WB 1:5000-1:50000 IF 1:50-1:500

Applications

Tested Applications: FC, IF/ICC, WB, ELISA Cited Applications: WB, IF, FC Species Specificity:

Human, mouse, rat **Cited Species:** human, rat, mouse

Positive Controls:

WB: HeLa cells, A431 cells, NIH/3T3 cells, rat liver tissue, HEEK-293 cells, Jurkat cells, HSC-T6 cells, RAW 264.7 cells, HEK-293 cells

IF: HepG2 cells,

Background Information

Ribosomal protein S6 (RPS6), Phosphoprotein NP33.1t may play an important role in controlling cell growth and proliferation through the selective translation of particular classes of mRNA. Ribosomal protein S6 is the major substrate of protein kinases in eukaryote ribosomes. The phosphorylation is stimulated by growth factors, tumor promoting agents, and mitogens. It is dephosphorylated at growth arrest. Phosphorylated at Ser-235 and Ser-236 by RPS6KA1 and RPS6KA3; phosphorylation at these sites facilitates the assembly of the preinitiation complex.

Notable Publications

Author	Pubmed ID	Journal	Application
Yifan Hong	34649140	Ecotoxicol Environ Saf	WB
Dongdong Yang	36417878	Cell Rep	WB
Yaoxiang Tang	35596155	BMC Cancer	WB

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66886-1-lg (56 Ribosomal protein antibody) at dilution of 1:500 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using 66886-1-1g (56 Ribosomal protein antibody) at dilution of 1:100 and CoraLite488-Conjugated Affini Pure Goat Anti-Mouse IgG(H+L).



Various lysates were subjected to SDS PAGE followed by western blot with 66886-1-lg (S6 Ribosomal protein antibody) at dilution of 1:15000 incubated at room temperature for 1.5 hours.



1X10^6 HepG2 cells were intracellularly stained with 0.4 ug Anti-Human S6 Ribosomal protein (66886-1-lg, Clone:1C3E10) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Control Antibody. Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).