For Research Use Only

Alpha Actinin Monoclonal antibody

Catalog Number:66895-1-lg 7 Publications



Basic Information

Catalog Number: 66895-1-Ig

Size: GeneID (NCBI): 1500 μ g/ml 87

Source: UNIPROT ID:
Mouse P12814
Isotype: Full Name:
IgG2b actinin, alpha 1
Immunogen Catalog Number: Calculated MW:
AG28383 103 kDa

Observed MW: 100-105 kDa

BC015766

GenBank Accession Number:

Purification Method:

Protein A purification CloneNo.:

1E2E1

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:2000 IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

Cited Applications:

WB, IF

Species Specificity: Human, rat, mouse Cited Species:

human, mouse, rat, mosquito

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

IF/ICC: C2C12 cells,

WB: HeLa cells, RAW 264.7 cells, HEK-293 cells, HepG2 cells, Jurkat cells, HSC-T6 cells

IHC: mouse heart tissue,

Background Information

Alpha Actinins are highly conserved actin-binding proteins that form antiparallel homodimers with the actin-binding domains on each end of the molecule, allowing crosslinking of actin molecules. There are four alpha actinin isoforms identified: two ubiquitously expressed cytoskeletal isoforms (ACTN1 and ACTN4) and two striated muscle-specific sarcomeric isoforms (ACTN2 and ACTN3). ACTN1 is also alternatively spliced to form smooth muscle or brain specific isoforms. This antibody recognize endogenous ACTN1, while it may cross-react with ACTN2 and ACTN4.

Notable Publications

Author	Pubmed ID	Journal	Application
Jialu Qiao	34826589	Cell Signal	WB
Zequn Zheng	35873575	Front Pharmacol	IF
Jiayu Ren	39158709	Cell Mol Life Sci	IF

Storage

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66895-1-Ig (a-Actinin antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 66895-1-Ig (alpha Actinin antibody) at dilution of 1:1000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded mouse heart tissue slide using 66895-1-Ig (alpha Actinin antibody) at dilution of 1:1000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10^6 A431 cells were intracellularly stained with 0.4 ug Anti-Human Alpha Actinin (66895-1-lg, Clone:1E2E1) and Coralite® 488-Conjugated Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2b Isotype Control (MPC-11) (65128-1-lg, Clone: MPC-11) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).



Immunofluorescent analysis of (-20°C Methanol) fixed C2C12 cells using Alpha Actinin antibody (66895-1-Ig, Clone: 1E2E1) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L).