For Research Use Only

SPTLC1 Monoclonal antibody

Catalog Number:66899-1-lg

Featured Product

1 Publications

BC007085

10558

015269

GeneID (NCBI):

UNIPROT ID:

Full Name:

chain base subunit 1

Calculated MW: 53 kDa Observed MW: 55 kDa

GenBank Accession Number:

serine palmitoyltransferase, long



Basic Information

Applications

Catalog Number: 66899-1-lg Size:

981 ug/ml Source: Mouse Isotype: lgG1

Immunogen Catalog Number:

AG7876

Tested Applications:

WB, IHC, IF/ICC, ELISA **Cited Applications:**

Species Specificity: human, mouse, rat Cited Species:

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Purification Method: Protein G purification

CloneNo.: 2D11D7

Recommended Dilutions: WB 1:5000-1:50000

IHC 1:50-1:500 IF/ICC 1:490-1:1962

Positive Controls:

WB: A549 cells, HEK-293 cells, RAW 264.7 cells, HT-29 cells, HeLa cells, HepG2 cells, Jurkat cells, K-562 cells,

HSC-T6 cells, NIH/3T3 cells IHC: human colon cancer tissue,

IF/ICC: HepG2 cells,

Background Information

SPTLC1 is a subunit of serine palmitoyltransferase (SPT) which is the key enzyme in sphingolipid biosynthesis and is essential for embryogenesis and cell survival. Mutations in the SPTLC1 gene (C133W, C133Y, V144D, and G387A) were reported to be responsible for the development of an inherited sensory neuropathy (hereditary sensory neuropathy type I, HSN1).

Notable Publications

Author **Pubmed ID** Journal Application Nucleic Acids Res WB Xianping Shi 33080033

Storage


Storage:

Store at -20°C. Stable for one year after shipment.

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66899-1-1g (SPTLC 1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 66899-1-Ig (SPTLC1 antibody) at dilution of 1:500 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 66899-1-Ig (SPTLC1 antibody) at dilution of 1:500 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



WB result of SPTLC1 antibody (66899-1-lg; 1:2000; incubated at room temperature for 1.5 hours) with sh-Control and sh-SPTLC1 transfected HEK-293 cells.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using SPTLC1 antibody (66899-1-lg, Clone: 2D11D7) at dilution of 1:981 and Multi-rAb CoraLite® Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (Cat.NO. RGAM004).