For Research Use Only

MAVS; VISA Monoclonal antibody

lgG1

Catalog Number:66911-1-lg Featured Product

12 Publications



Basic Information

Catalog Number: GenBank Accession Number: 66911-1-lg BC044952 GeneID (NCBI): Concentration: 2524 ug/ml 57506 **UNIPROT ID:** Source: Mouse Q7Z434 Full Name: Isotype:

mitochondrial antiviral signaling

protein Immunogen Catalog Number:

AG5949 Calculated MW:

57 kDa Observed MW: 50-55 kDa, 70-75 kDa

Applications

Tested Applications: WB, IHC, ELISA **Cited Applications:** WB, IF, IP, CoIP Species Specificity: human

Cited Species:

human, mouse, pig, monkey

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Purification Method:

Protein A purification

CloneNo.: 1A8E9

Recommended Dilutions:

WB 1:5000-1:50000 IHC 1:550-1:2200

Positive Controls:

WB: A431 cells, HeLa cells, HepG2 cells, Jurkat cells,

LO2 cells, HEK-293 cells, THP-1cells

IHC: human liver cancer tissue.

Background Information

 $Mit ochondrial\ antiviral\ - signaling\ protein\ (MAVS)\ is\ also\ known\ as\ virus\ - induced\ - signaling\ adapter\ (VISA)\ or\ IFN$ beta promoter stimulator protein 1 (IPS-1), it is widely involved and required for innate immune defense against viruses. MAVS, present in T cells, monocytes, epithelial cells and hepatocytes, contains CARD and transmembrane domains which are essential for antiviral functions. MAVS is able to interact with various cellular proteins including DDX58/RIG-I, IFIH1/MDA5, TRAF2, TRAF6, TMEM173/MITA, IFIT3 and etc. It can undergoe phosphorylation on multiple sites and ubiquitination, which may together cause the molecular weight migrate to about 70 kDa despite the predicated 57 kDa.

Notable Publications

Author	Pubmed ID	Journal	Application
Zhihai Zhou	33692778	Front Immunol	WB
Xiaohua Jie	35121645	J Immunother Cancer	WB
Yumei Han	33328314	J Virol	WB

Storage

Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




A431 cells were subjected to SDS PAGE followed by western blot with 66911-1-1g (MAVS; VISA antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 66911-1-1g (MAVS; VISA antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 66911-1-Ig (MAVS; VISA antibody) at dilution of 1:1100 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 66911-1-Ig (MAVS; VISA antibody) at dilution of 1:1100 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



WB result of MAVS; VISA antibody (66911-1-lg; 1:4000; incubated at room temperature for 1.5 hours) with sh-Control and sh-MAVS; VISA transfected HeLa cells.