For Research Use Only

## ROR1 Monoclonal antibody, PBS Only (Detector)

Catalog Number:66923-1-PBS

Featured Product




Basic Information	Catalog Number: 66923-1-PBS	GenBank Accession Number: BC006374	Purification Method: Protein A purification	
	Size: 1mg/ml	GenelD (NCBI): 4919	CloneNo.: 1F5D8	
	Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG14586	UNIPROT ID: Q01973		
		Full Name: receptor tyrosine kinase-like orphan receptor 1 Calculated MW: 937 aa, 104 kDa		
				Observed MW: 130 kDa
		Applications	Tested Applications: WB, IHC, Cytometric bead array, I Species Specificity: human, mouse	WB, IHC, Cytometric bead array, Indirect ELISA Species Specificity:
Background Informatic	ROR1 is a member of the RTK family of orphan receptors related to muscle-specific kinase and Trk neurotrophin receptors (PMID: 18546292). ROR1 is primarily expressed by neural crest cells during embryogenesis. High expression of ROR1 is reported in many types of malignancies and is thought to be involved in tumor growth, apoptosis, and epithelial-mesenchymal transition (PMID: 26245996). The human ROR1 gene encodes a 937 amino acid protein with an Ig-like domain, a cysteine-rich domain, a kringle domain, a tyrosine kinase domain, and a proline-rich domain. The calculated molecular weight of ROR1 is 104 kDa, but ROR1 has multiple N-glycosylation sites that generate post-translationally modified ROR1 at 130 kDa (PMID: 24752542).			
Storage	Storage: Store at -80°C. The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer: PBS Only			

For technical support and original validation data for this product please contact:T: 4006900926E: Proteintech-CN@ptglab.comW: ptgcn.com


This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66923-1-lg (ROR1 antibody) at dilution of 1:3000 incubated at room

antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control. This data was developed using the same antibody clone with 66923-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66923-1-1g (ROR1 antibody) at dilution of 1:600 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66923-1-PBS in a different storage buffer formulation.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 66923-1-Ig (ROR1 antibody) at dilution of 1:600 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 66923-1-PBS in a different storage buffer formulation.



NIH/3T3 cells were subjected to SDS PAGE followed by western blot with 66923-1-lg (ROR1 antibody) at dilution of 1:3000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 66923-1-PBS in a different storage buffer formulation.



Cytometric bead array standard curve of MP51139-1, ROR1 Monoclonal Matched Antibody Pair, PBS Only. Capture antibody: 66923-2-PBS. Detection antibody: 66923-1-PBS. Standard:Ag14586. Range: 0.391-100 ng/mL



Cytometric bead array standard curve of MP51139-2, ROR1 Monoclonal Matched Antibody Pair, PBS Only. Capture antibody: 66923-3-PBS. Detection antibody: 66923-1-PBS. Standard:Ag14586. Range: 0.781-100 ng/mL