

For Research Use Only

DDX21 Monoclonal antibody

Catalog Number: 66925-1-Ig **2 Publications**



Basic Information

Catalog Number: 66925-1-Ig	GenBank Accession Number: BC008071	Purification Method: Protein A purification
Size: 1300 ug/ml	GeneID (NCBI): 9188	CloneNo.: 1B9C8
Source: Mouse	UNIPROT ID: Q9NR30	Recommended Dilutions: WB 1:2000-1:10000 IHC 1:250-1:1000 IF/ICC 1:400-1:1600
Isotype: IgG1	Full Name: DEAD (Asp-Glu-Ala-Asp) box polypeptide 21	
Immunogen Catalog Number: AG17850	Calculated MW: 783 aa, 87 kDa	
	Observed MW: 90-95 kDa	

Applications

Tested Applications: WB, IHC, IF/ICC, FC (Intra), ELISA	Positive Controls:
Cited Applications: WB, IHC, IP	WB : HeLa cells, HEK-293 cells, HepG2 cells, Jurkat cells, K-562 cells, HL-60 cells, THP-1 cells
Species Specificity: human	IHC : human colon cancer tissue, human colon tissue
Cited Species: human	IF/ICC : HepG2 cells,
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

DDX21 protein belongs to DEAD box protein family which is characterized by the conserved motif Asp-Glu-Ala-Asp (DEAD). As a putative RNA helicase, DDX21 unwinds double-stranded RNA, folds single-stranded RNA and is involved in process including ribosomal RNA biogenesis, RNA editing and general transcription. Interaction of DDX21 and c-Jun was reported in ribosomal RNA processing.

Notable Publications

Author	Pubmed ID	Journal	Application
Katherine L Harper	39592606	Nat Commun	
Guanghui Zhang	39187490	Cell Death Dis	WB,IP,IHC

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 66925-1-Ig (DDX21 antibody) at dilution of 1:5000 incubated at room temperature for 1.5 hours.




Various lysates were subjected to SDS PAGE followed by western blot with 66925-1-Ig (DDX21 antibody) at dilution of 1:50000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Alpha Tubulin Monoclonal antibody (HRP-66031) as loading control.



Immunohistochemical analysis of paraffin-embedded human colon cancer tissue slide using 66925-1-Ig (DDX21 antibody) at dilution of 1:500 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using DDX21 antibody (66925-1-Ig, Clone: 1B9C8) at dilution of 1:800 and CoraLite@488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1), CL594-Phalloidin (red).



1X10⁶ HepG2 cells were intracellularly stained with 0.2 ug Anti-Human DDX21 (66925-1-Ig, Clone:1B9C8) and CoraLite@488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.2 ug Mouse IgG1 Isotype Control (66360-1-Ig, Clone: T1F8D3F10) (blue). Cells were fixed and permeabilized with Transcription Factor Staining Buffer Kit (PF00011).