#### For Research Use Only

# FGFR3 Monoclonal antibody

Catalog Number:66954-1-lg Featured Product

10 Publications



**Basic Information** 

Catalog Number: 66954-1-lg

GenBank Accession Number: NM 000142 GeneID (NCBI):

**Purification Method:** Protein G purification

Size: 1500 µg/ml

2261 Full Name:

Calculated MW:

CloneNo.: 1F3G1

Source: Mouse Isotype:

fibroblast growth factor receptor 3

Recommended Dilutions: WB 1:5000-1:50000

lgG1

87 kDa

Immunogen Catalog Number:

Observed MW: AG26290 125-135 kDa

IF/ICC 1:200-1:800

**Applications** 

**Tested Applications:** WB, IF/ICC, ELISA Cited Applications: WB, IHC, IF Species Specificity:

human **Cited Species:** human, mouse Positive Controls:

WB: LNCaP cells, L02 cells, L-929 cells, HeLa cells, HEK-293 cells, HepG2 cells, A549 cells, NCI-H1299 cells

IF/ICC: HepG2 cells,

## **Background Information**

Fibroblast growth factors (FGFs) are polypeptide growth factors involved in a variety of activities including mitogenesis, angiogenesis, and wound healing (PMID: 1847508). The human FGF receptor family, a subfamily of receptor tyrosine kinases (RTKs), comprises of four family members-FGFR1, FGFR2, FGFR3, and FGFR4 (PMID: 23900974). Each receptor contains an extracellular domain with either two or three immunoglobulin-like domains, a transmembrane domain, and a cytoplasmic tyrosine kinase domain. FGFR3 binds acidic and basic fibroblast GH and plays a role in bone development and maintenance. Mutations in the FGFR3 gene lead to craniosynostosis and multiple types of skeletal dysplasia. Due to frequent mutations in certain cancers, the FGFR3 gene has also been associated with tumor progression.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Liang Kuang	31662319	Ann Rheum Dis	WB
Limin Wang	36305369	Tissue Eng Part A	IF
Fake Liao	34787070	Bioengineered	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


### **Selected Validation Data**




Various lysates were subjected to SDS PAGE followed by western blot with 66954-1-1g (FGFR3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using FGFR3 antibody (66954-1-lg, Clone: 1F3G1) at dilution of 1:400 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using FGFR3 antibody (66954-1-lg, Clone: 1F3G1) at dilution of 1:400 and CoraLite® 488-Conjugated Goat Anti-Mouse IgG(H+L).



L-929 cells were subjected to SDS PAGE followed by western blot with 66954-1-1g (FGFR3 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.