

For Research Use Only

RRM2 Monoclonal antibody

Catalog Number: 67006-1-Ig **4 Publications**



Basic Information

Catalog Number:

67006-1-Ig

Size:

1000 ug/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG28664

GenBank Accession Number:

BC030154

GeneID (NCBI):

6241

UNIPROT ID:

P31350

Full Name:

ribonucleotide reductase M2 polypeptide

Calculated MW:

389 aa, 45 kDa

Observed MW:

45 kDa

Purification Method:

Protein A purification

CloneNo.:

2A9A7

Recommended Dilutions:

WB 1:5000-1:50000

IP 0.5-4.0 ug for 1.0-3.0 mg of total protein lysate

IHC 1:2000-1:8000

IF/ICC 1:200-1:800

Applications

Tested Applications:

WB, IHC, IF/ICC, IP, ELISA

Cited Applications:

WB, IF, CoIP

Species Specificity:

human

Cited Species:

human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB : HeLa cells, HEK-293 cells, MCF-7 cells, A431 cells, Jurkat cells, K-562 cells, THP-1 cells

IP : HEK-293 cells,

IHC : human skin cancer tissue,

IF/ICC : HepG2 cells,

Background Information

Ribonucleotide reductase M2 subunit is one of two subunits that constitute ribonucleotide reductase, the enzyme that catalyzes the conversion of ribonucleotide 5'-diphosphates into 2'-deoxyribonucleotides, a rate-limiting step in the production of 2'-deoxyribonucleoside 5'-diphosphates (dNTP) required for DNA synthesis and repair that is required for DNA synthesis and repair [PMID:20825972, 19250552]. RRM2 is only expressed during the late G1/early S phase, and degraded in late S phase, and the activity of RNR, and therefore DNA synthesis and cell proliferation, is controlled during the cell cycle by the synthesis and degradation of RRM2 subunit [PMID:3894352].

Notable Publications

Author	Pubmed ID	Journal	Application
S M Du	32122142	Neoplasma	WB
Heng Gao	39630361	Mol Cell Biochem	WB
Zhouyuan Du	39398252	iScience	IF

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67006-1-Ig (RRM2 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



IP result of anti-RRM2 (IP:67006-1-Ig, 4ug; Detection:67006-1-Ig 1:2000) with HEK-293 cells lysate 1470 ug.



Immunohistochemical analysis of paraffin-embedded skin cancer slide using 67006-1-Ig (RRM2 antibody) at dilution of 1:4000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Ethanol) fixed HepG2 cells using RRM2 antibody (67006-1-Ig, Clone: 2A9A7) at dilution of 1:400 and Multi-rAb CoraLite ® Plus 488-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM002).