

For Research Use Only

# ESD Monoclonal antibody

Catalog Number: 67069-2-Ig



## Basic Information

Catalog Number:

67069-2-Ig

Size:

1000 ug/ml

Source:

Mouse

Isotype:

IgG1

Immunogen Catalog Number:

AG7487

GenBank Accession Number:

BC001169

GeneID (NCBI):

2098

UNIPROT ID:

P10768

Full Name:

esterase D/formylglutathione  
hydrolase

Calculated MW:

31 kDa

Observed MW:

30-34 kDa

Purification Method:

Protein G purification

CloneNo.:

2F5E1

Recommended Dilutions:

WB 1:5000-1:50000

IF/ICC 1:200-1:800

## Applications

Tested Applications:

WB, IF/ICC, ELISA

Species Specificity:

human

Positive Controls:

WB : HCT 116 cells, Caco-2 cells, HepG2 cells, Jurkat  
cells, K-562 cells, Ramos cells

IF/ICC : HEK-293 cells,

## Background Information

Esterase D (ESD) is a non-specific esterase widely distributed in various organisms and is also named S-Formylglutathione Hydrolase (SFGH). ESD is a member of the carboxylesterase family and has both carboxylesterase and thioesterase activities. ESD plays an important role in the process of glutathione-dependent detoxification, regulating cholesterol efflux and virus infection in humans, and is closely related to the development of tumors. ESD as a Genetic Marker for Retinoblastoma (PMID: 32247735, PMID: 34875997, PMID: 35627173). The calculated molecular weight of ESD is 31 kDa.

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67069-2-Ig (ESD antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Ethanol) fixed HEK-293 cells using ESD antibody (67069-2-Ig, Clone: 2F5E1) at dilution of 1:400 and Multi-rAb CoraLite® Plus 488-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM002).