

For Research Use Only

# PI3 Kinase p110 Alpha Monoclonal antibody

Catalog Number: 67071-1-Ig

Featured Product

118 Publications



## Basic Information

Catalog Number:

67071-1-Ig

Concentration:

1000 ug/ml

Source:

Mouse

Isotype:

IgG2b

Immunogen Catalog Number:

AG17392

GenBank Accession Number:

BC113601

GeneID (NCBI):

5290

UNIPROT ID:

P42336

Full Name:

phosphoinositide-3-kinase, catalytic, alpha polypeptide

Calculated MW:

1068 aa, 124 kDa

Observed MW:

110 kDa

Purification Method:

Protein A purification

CloneNo.:

1F6A7

Recommended Dilutions:

WB: 1:1000-1:4000

IHC: 1:250-1:1000

IF/ICC: 1:200-1:800

FC (Intra): 0.40 ug per 10<sup>6</sup> cells in a 100 µl suspension

## Applications

Tested Applications:

WB, IHC, IF/ICC, FC (Intra), ELISA

Cited Applications:

WB, IHC, IP, CoIP, RIP

Species Specificity:

human, mouse, rat, pig, rabbit

Cited Species:

human, mouse, rat, monkey, bovine, fish

**Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0**

Positive Controls:

WB: U2OS cells, pig brain tissue, LNCaP cells, HeLa cells, HEK-293 cells, Jurkat cells, K-562 cells, rabbit brain tissue, rat brain tissue, mouse brain tissue

IHC: human breast cancer tissue,

IF/ICC: HeLa cells,

FC (Intra): Jurkat cells,

## Background Information

PIK3CA belongs to the PI3/PI4-kinase family. It phosphorylates PtdIns, PtdIns4P and PtdIns(4,5)P<sub>2</sub> with a preference for PtdIns(4,5)P<sub>2</sub>. Defects in PIK3CA are associated with colorectal cancer (CRC). Defects in PIK3CA are associated with breast cancer. Defects in PIK3CA are associated with ovarian cancer. Defects in PIK3CA may underlie hepatocellular carcinoma (HCC). Defects in PIK3CA are a cause of keratosis seborrheic (KERSEB).

## Notable Publications

Author	Pubmed ID	Journal	Application
Lei Zhang	34592228	Life Sci	WB
Bailu Duan	36073658	Chem Biodivers	WB
Yujing Sun	32965543	J Cancer Res Clin Oncol	WB

## Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: [Proteintech-CN@ptglab.com](mailto:Proteintech-CN@ptglab.com)


W: [ptgcn.com](http://ptgcn.com)

**This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.**


## Selected Validation Data




U2OS cells were subjected to SDS PAGE followed by western blot with 67071-1-Ig (PI3 Kinase p110 Alpha antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. The membrane was stripped and reblotted with HRP-conjugated Lamin B1 Monoclonal antibody (HRP-66095) as loading control.




Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 67071-1-Ig (PIK3CA antibody) at dilution of 1:500 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human breast cancer tissue slide using 67071-1-Ig (PIK3CA antibody) at dilution of 1:500 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (-20°C Methanol) fixed HeLa cells using PI3 Kinase p110 Alpha antibody (67071-1-Ig, Clone: 1F6A7) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



1X10<sup>6</sup> Jurkat cells were intracellularly stained with 0.4 ug Anti-Human PI3 Kinase p110 Alpha (67071-1-Ig, Clone:1F6A7) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2b Isotype Control (MPC-11) (65128-1-Ig, Clone: MPC-11) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).