## For Research Use Only

# Endoglin/CD105 Monoclonal antibody

Catalog Number:67075-1-lg 7 Publications



**Basic Information** 

Catalog Number: 67075-1-lg Concentration: 1000 ug/ml

Source: Mouse Isotype: lgG2b Immunogen Catalog Number:

AG27837

Observed MW:

95-100 kDa

BC014271

2022

P17813

GeneID (NCBI):

**UNIPROT ID:** 

Full Name:

endoglin Calculated MW:

71 kDa

GenBank Accession Number:

**Purification Method:** 

Protein A purification CloneNo.:

2D5E8

Recommended Dilutions: WB 1:5000-1:20000 IHC 1:10000-1:40000 IF-P 1:5000-1:20000

**Applications** 

**Tested Applications:** WB, IHC, IF-P, ELISA **Cited Applications:** IHC, IF

Species Specificity: human

**Cited Species:** 

human, mouse, rat, sheep

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (\*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

#### Positive Controls:

WB: HL-60 cells, HUVEC cells, human placenta tissue

IHC: human breast cancer tissue, human liver cancer tissue, human placenta tissue, human renal cell carcinoma tissue, human tonsillitis tissue

IF-P: human tonsillitis tissue, human liver cancer tissue, human placenta tissue

## **Background Information**

Endoglin (ENG, CD105) is a homodimeric cell membrane glycoprotein of 180 kDa, composed of disulphide-linked subunits of 95 kDa. Endoglin is a proliferation-associated and hypoxia-inducible protein mainly expressed on vascular endothelial cells. It acts as an accessory receptor for transforming growth factor beta (TFG-  $\beta$  ) and is involved in vascular development and remodelling. The important role of Endoglin in angiogenesis and in tumor progression makes it an ideal target for antiangiogenic therapy and a good marker for tumor prognosis. (PMID: 1326540; 14528280; 21737653; 12773481)

## **Notable Publications**

Author	Pubmed ID	Journal	Application
Qing Zhang	35765095	Stem Cell Res Ther	IF
Baihe Han	35968497	Dis Markers	FC
Hou-Fu Xia	35916917	Cell Tissue Res	IHC,IF

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol, pH7.3


Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926 E: Proteintech-CN@ptglab.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67075-1-1g (Endoglin/CD105 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 67075-1-lg (Endoglin/CD105 antibody) at dilution of 1:20000 (under 10x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human breast cancer tissue slide using 67075-1-lg (Endoglin/CD105 antibody) at dilution of 1:20000 (under 40x lens. Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using Endoglin/CD105 antibody (67075-1-lg, Clone: 2D5E8) at dilution of 1:10000 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using Endoglin/CD105 antibody (67075-1-lg, Clone: 2D5E8) at dilution of 1:10000 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunohistochemical analysis of paraffinembedded human liver cancer tissue slide using 67075-1-lg (Endoglin/CD105 antibody) at dilution of 1:20000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed paraffin-embedded human placenta tissue using Endoglin/CD105 antibody (67075-1-lg, Clone: 2D5E8) at dilution of 1:400 and Coralite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).