

EAAT2 Monoclonal antibody

Catalog Number: 67083-1-Ig 1 Publications

Basic Information

Catalog Number: 67083-1-Ig	GenBank Accession Number: BC132768	Purification Method: Protein A purification
Size: 1400 µg/ml	GeneID (NCBI): 6506	CloneNo.: 1C8D10
Source: Mouse	UNIPROT ID: P43004	Recommended Dilutions: WB 1:1000-1:8000 IHC 1:200-1:800
Isotype: IgG1	Full Name: solute carrier family 1 (glial high affinity glutamate transporter), member 2	
Immunogen Catalog Number: AG19816	Calculated MW: 574 aa, 62 kDa	
	Observed MW: 60-70 kDa	

Applications

Tested Applications: IHC, WB, ELISA	Positive Controls:
Cited Applications: WB	WB : rat brain tissue, rat cerebellum tissue, mouse brain tissue
Species Specificity: Human, mouse, rat	IHC : mouse brain tissue, rat brain tissue
Cited Species: mouse	
Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0	

Background Information

EAAT2 (also known as GLT-1), is encoded by SLC1A2 and is a glutamate transporter which can clear the majority of extracellular glutamate in brain and prevent brain seizures. Three isoforms of EAAT2 exist due to the alternative splicing. In addition to the 65-70 kDa monomer form, 130-150 kDa dimer of EAAT2 can also be observed on SDS-PAGE. (PMID: 22114258)


Notable Publications

Author	Pubmed ID	Journal	Application
Jing Yan	38325382	Cell Rep Med	WB




Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67083-1-Ig (SLC1A2 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded rat brain tissue slide using 67083-1-Ig (EAAT2 antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded mouse brain tissue slide using 67083-1-Ig (EAAT2 antibody) at dilution of 1:400 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).