For Research Use Only

L1CAM Monoclonal antibody

Catalog Number:67115-1-lg 3 Publications



Basic Information

Catalog Number: 67115-1-lg Size:

1000 ug/ml
Source:
Mouse
Isotype:
IgG1

Immunogen Catalog Number:

AG17706

Observed MW:

220 kDa

BC126229

3897

P32004

GeneID (NCBI):

UNIPROT ID:

Full Name:

1257 aa, 140 kDa

GenBank Accession Number:

L1 cell adhesion molecule Calculated MW: Purification Method:

Protein A purification CloneNo.: 3H7B9

Recommended Dilutions: WB 1:5000-1:50000 IHC 1:500-1:2000 IF-P 1:200-1:800

Applications

Tested Applications: WB, IHC, IF-P, ELISA Cited Applications: WB. IF

Species Specificity:

human, mouse, rat, pig, rabbit

Cited Species: human, mouse

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: pig brain tissue, pig cerebellum tissue, Rabbit brain tissue, rat brain tissue, mouse brain tissue

IHC: human colon cancer tissue, human brain tissue, human colon tissue, mouse brain tissue

IF-P: mouse brain tissue,

Background Information

L1CAM, also known as NCAM-L1 or CD171, is a cell adhesion molecule of the immunoglobulin superfamily. It is a 200-220 kDa transmembrane glycoprotein composed of six Ig-like domains and five fibronectin type III repeats followed by a transmembrane region and a highly conserved cytoplasmic tail (PMID: 3412448; 22796939). L1CAM is primarily expressed in the nervous system and is involved in neuron-neuron adhesion, neurite fasciculation, outgrowth of neurites, cerebellar granule cell migration, neurite outgrowth on Schwann cells and interactions among epithelial cells of intestinal crypts (PMID: 3412448; 10767310). L1CAM is overexpressed in many human cancers and is often associated with bad prognosis (PMID: 27267927; 26111503).

Notable Publications

Author	Pubmed ID	Journal	Application
Danyu Li	38898558	J Extracell Vesicles	WB
Xinli Jiang	38647743	J Neuroimmune Pharmacol	WB
Yudong Liu	38424563	Cell Commun Signal	WB,IF

Storage

Storage

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:

T: 4006900926 E: Proteintech-CN@ptglab.com

W: ptgcn.co

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67115-1-lg (L1CAM antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.




Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 67115-1-Ig (L1CAM antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human colon cancer tissue slide using 67115-1-Ig (L1CAM antibody) at dilution of 1:1000 (under 20x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using L1CAM antibody (67115-1-lg, Clone: 3H7B9) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed mouse brain tissue using L1CAM antibody (67115-1-Ig, Clone: 3H7B9) at dilution of 1:400 and CoraLite® 488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).

Immunohistochemical analysis of paraffinembedded human colon tissue slide using 67115-1-Ig (L1CAM antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).