

For Research Use Only

RANGAP1 Monoclonal antibody

Catalog Number: 67146-1-Ig



Basic Information

Catalog Number: 67146-1-Ig	GenBank Accession Number: BC014044	Purification Method: Protein A purification
Size: 2000 µg/ml	GeneID (NCBI): 5905	CloneNo.: 1F3A4
Source: Mouse	UNIPROT ID: P46060	Recommended Dilutions: WB 1:5000-1:50000 IF 1:50-1:200
Isotype: IgG1	Full Name: Ran GTPase activating protein 1	
Immunogen Catalog Number: AG26136	Calculated MW: 64 kDa	
	Observed MW: 64 kDa, 82 kDa	

Applications

Tested Applications: IF, WB, ELISA	Positive Controls:
Species Specificity: Human, mouse, rat	WB : U2OS cells, HeLa cells, HEK-293 cells, MCF-7 cells, Jurkat cells, HSC-T6 cells, NIH/3T3 cells, 4T1 cells, A549 cells, LNCaP cells, K-562 cells
	IF : HepG2 cells,

Background Information

RanGAP1 is a protein that associates with the nuclear pore complex and participates in the regulation of nuclear transport. In mammalian cells, unmodified RanGAP1 is predominantly cytoplasmic, whereas modification by small ubiquitin-related modifier protein (SUMO) targets RanGAP1 to the cytoplasmic filaments of nuclear pore complex (NPC) where it forms a complex with RanBP2 and Ubc9. Phosphorylation of RanGAP1 occurs in a cell-cycle-dependent manner and may play a role in regulating RanGAP1 catalytic activity. We got 64 kDa (cytoplasmic RanGAP1) and 82 kDa (SUMO-1 modified Ran GAP1) in western blotting.

Storage

Storage:
Store at -20°C. Stable for one year after shipment.
Storage Buffer:
PBS with 0.02% sodium azide and 50% glycerol pH 7.3.
Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



HeLa cells were subjected to SDS PAGE followed by western blot with 67146-1-Ig (RANGAP1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Various lysates were subjected to SDS PAGE followed by western blot with 67146-1-Ig (RANGAP1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (4% PFA) fixed HepG2 cells using 67146-1-Ig (RANGAP1 antibody) at dilution of 1:100 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).