For Research Use Only

CD138/Syndecan-1 Monoclonal antibody

Proteintech®

Antibodies | ELISA kits | Proteins

www.ptglab.com

Catalog Number:67155-1-lg 3 Publications

Basic Information

Catalog Number: 67155-1-lg

Size: 2108 ug/ml

Source:

BC008765 GeneID (NCBI): 6382

UNIPROT ID:

GenBank Accession Number:

MouseP18827Isotype:Full Name:IgG2asyndecan 1Immunogen Catalog Number:Calculated MW:AG082332.5 kDa

Purification Method:

Protein A purification

CloneNo.: 4F3A8

Recommended Dilutions: IHC 1:4000-1:16000 IF-P 1:200-1:800

Applications

Tested Applications: IHC, IF-P, ELISA Cited Applications:

IF

Species Specificity:

human
Cited Species:
human

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

IHC: human tonsillitis tissue, human lymphoma tissue, human ovary tumor tissue

IF-P: human tonsillitis tissue,

Background Information

Syndecan-1 (SDC1), also named CD138, is an integral membrane protein. It participates in cell proliferation, cell migration and cell-matrix interactions via its receptor for extracellular matrix proteins. It is a heparan sulfate proteoglycan expressed on the surface of, and actively shed by, myeloma cells. Altered syndecan-1 expression has been detected in several different tumor types. SDC1 was regarded as a useful marker for labeling normal and neoplastic plasma cells and plasmacytoid lymphomas.

Notable Publications

Author	Pubmed ID	Journal	Application
Yanglin Chen	38582082	Dev Cell	IF
Pei-Fen Zhang	36746966	Nat Commun	IF
Wang Zheng	35913916	Autophagy	IF

Storage

Storage:


Storage Ruffer

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 67155-1-Ig (CD138/Syndecan-1 antibody) at dilution of 1:4000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).




Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 67155-1-Ig (SDC1,CD138 antibody) at dilution of 1:8000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffinembedded human tonsillitis tissue slide using 67155-1-Ig (SDC1,CD138 antibody) at dilution of 1:8000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD138/Syndecan-1 antibody (67155-1-Ig, Clone: 4F3A8) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).



Immunofluorescent analysis of (4% PFA) fixed human tonsillitis tissue using CD138/Syndecan-1 antibody (67155-1-Ig, Clone: 4F3A8) at dilution of 1:400 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L).