

For Research Use Only

ECD Monoclonal antibody

Catalog Number: 67179-1-Ig



Basic Information

Catalog Number: 67179-1-Ig	GenBank Accession Number: BC000721	Purification Method: Protein G purification
Size: 800 µg/ml	GeneID (NCBI): 11319	CloneNo.: 1D5E8
Source: Mouse	UNIPROT ID: O95905	Recommended Dilutions: WB 1:20000-1:100000 IHC 1:500-1:2000
Isotype: IgG1	Full Name: ecdysoneless homolog (Drosophila)	
Immunogen Catalog Number: AG28674	Calculated MW: 73 kDa	
	Observed MW: 70-73 kDa	

Applications

Tested Applications:

IHC, WB, ELISA

Species Specificity:

Human, mouse, rat

Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0

Positive Controls:

WB: HeLa cells, NIH/3T3 cells, HSC-T6 cells, Jurkat cells, JAR cells, NCCIT cells, HEK-293 cells

IHC: human renal cell carcinoma tissue, human stomach cancer tissue, mouse bladder tissue, rat colon tissue, rat bladder tissue

Background Information

Storage

Storage:

Store at -20°C. Stable for one year after shipment.

Storage Buffer:

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

For technical support and original validation data for this product please contact:


T: 4006900926

E: Proteintech-CN@ptglab.com


W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67179-1-Ig (ECD antibody) at dilution of 1:100000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human renal cell carcinoma tissue slide using 67179-1-Ig (ECD antibody) at dilution of 1:1000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human renal cell carcinoma tissue slide using 67179-1-Ig (ECD antibody) at dilution of 1:1000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).