For Research Use Only

ENO1 Monoclonal antibody

Catalog Number:67187-1-lg Featured Product



Basic Information

Catalog Number:

GenBank Accession Number:

Purification Method:

67187-1-lg Size: BC015641 GeneID (NCBI): Thiophilic affinity chromatograph

200 ug/ml Source: GeneID (NCBI): 2023 CloneNo.: 1D6E8

Mouse Isotype: ENSEMBL Gene ID: ENSG00000074800 UNIPROT ID: Recommended Dilutions: WB 1:5000-1:50000 IF/ICC 1:400-1:1600

Immunogen Catalog Number: AG17518

enolase 1, (alpha)

P06733

Full Name:

Calculated MW: 47 kDa Observed MW: 47 kDa

Applications

Tested Applications:

WB, IF/ICC, ELISA

Species Specificity:
human, mouse, rat, pig

Positive Controls:

WB: HEK-293 cells, HeLa cells, PC-3 cells, Jurkat cells, pig brain tissue, rat brain tissue, mouse brain tissue

IF/ICC: HeLa cells,

Background Information

ENO1, also named as NNE, ENO1L1, MBPB1, MPB1 and MBP1, belongs to the enolase family. ENO1 is a metabolic enzyme involved in the synthesis of pyruvate. It also acts as a plasminogen receptor and mediates the activation of plasmin and extracellular matrix degradation. In tumor cells, ENO1 is up-regulated and supports the Warburg effect; it is expressed at the cell surface, where it promotes cancer invasion, and is subjected to a specific array of post-translational modifications, namely acetylation, methylation and phosphorylation. ENO1 overexpression and post-translational modifications could be of diagnostic and prognostic value in many cancer types. (PMID: 27814656)

Storage

Storage:


Store at -20°C. Stable for one year after shipment.

Storage Buffer:


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67187-1-1g (ENO 1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



WB result of ENO1 antibody (67187-1-lg; 1:5000; incubated at room temperature for 1.5 hours) with sh-Control and sh-ENO1 transfected HEK-293 cells.



Immunofluorescent analysis of (-20°C Ethanol) fixed Hela cells using ENO1 antibody (67187-1-lg, Clone: 1D6E8) at dilution of 1:800 and CoraLite®488-Conjugated Goat Anti-Mouse IgG(H+L) (SA00013-1).