For Research Use Only

## TBK1 Monoclonal antibody, PBS Only

Catalog Number: 67211-1-PBS

**Featured Product** 



**Basic Information** 

Catalog Number:

GenBank Accession Number:

Purification Method: Protein A purification

67211-1-PBS

BC034950 GeneID (NCBI):

otein A punncatio

Size: 1 mg/ml

29110

CloneNo.: 2D7B1

Source: Mouse

UNIPROT ID: Q9UHD2 Full Name:

Isotype: IgG2b

TANK-binding kinase 1

Immunogen Catalog Number: AG28894

C Observed MW: 84 kDa

**Applications** 

Tested Applications:

WB,ELISA

Species Specificity:

Human, Mouse, Rat

## **Background Information**

TBK1, also named as tumor necrosis factor (TNF) receptor-associated factor NF-kB activator (TANK)-binding kinase 1 (TBK1), NF-kB-activating kinase (NAK), T2K, is a multimeric kinase that modulates inflammation and autophagy. It is a ubiquitously expressed serine-threonine kinase belonging to the 'noncanonical IkB kinases' (IKKs) recognized for its critical role in regulating type I IFN production (PMID: 27211305). And TBK1 is an important player in yet another critical cellular function, autophagy. This antibody is specific for TBK1.

Storage


Storage:

Store at -20  $^{\circ}\text{C}$  . Stable for one year after shipment.


Storage Buffer:

Aliquoting is unnecessary for -20°C storage


## Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67211-1-lg (TBK1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67211-1-PBS in a different storage buffer formulation.




Various lysates were subjected to SDS PAGE followed by western blot with 67211-1-1g (TBK1 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67211-1-PBS in a different storage buffer formulation.



U2OS (WT and TBK1 KO) lysates prepared with RIPA buffer, 50 µ g protein loaded. 67211-1-lg incubated at 1:5000 at 4°C overnight in 5% BSA in TBST. Ponceau stained transfers shown on right. Data provided by YCharOS, an open science company with a mission to validate commercial antibodies to improve scientific reproducibility and transparency. This data was developed using the same antibody clonewith 67211-1-PBS in a different storage buffer formulation. 

■ The provided Head of the same and the



Various lysates were subjected to SDS PAGE followed by western blot with 67211-1-lg (TBK1 antibody) at dilution of 1:4000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67211-1-PBS in a different storage buffer formulation.