

For Research Use Only

# NQO1 Monoclonal antibody

Catalog Number: 67240-1-Ig **110 Publications**



## Basic Information

Catalog Number: 67240-1-Ig	GenBank Accession Number: BC007659	Purification Method: Protein A purification
Concentration: 1000 ug/ml	GeneID (NCBI): 1728	CloneNo.: 1E5G7
Source: Mouse	UNIPROT ID: P15559	Recommended Dilutions: WB 1:5000-1:50000 IHC 1:2500-1:10000
Isotype: IgG2a	Full Name: NAD(P)H dehydrogenase, quinone 1	
Immunogen Catalog Number: AG28933	Calculated MW: 274 aa, 31 kDa	
	Observed MW: 29-31 kDa	

## Applications

Tested Applications: WB, IHC, FC (Intra), ELISA	Positive Controls: WB : HepG2 cells, HSC-T6 cells, L02 cells, K-562 cells IHC : human colon tissue,
Cited Applications: WB, IHC, IF	
Species Specificity: human, rat	
Cited Species: human, mouse, rat, pig	
<b>Note-IHC: suggested antigen retrieval with TE buffer pH 9.0; (*) Alternatively, antigen retrieval may be performed with citrate buffer pH 6.0</b>	

## Background Information

NQO1, also named as DIA4, NMOR1, DTD and QR1, belongs to the NAD(P)H dehydrogenase (quinone) family. This enzyme apparently serves as a quinone reductase in connection with conjugation reactions of hydroquinones involved in detoxification pathways as well as in biosynthetic processes such as the vitamin K-dependent gamma-carboxylation of glutamate residues in prothrombin synthesis. It is known to be involved in benzene metabolism. In human studies of ozone exposure, polymorphisms in oxidative stress genes (NQO1, GSTM1, GSTP1) modify respiratory symptoms, lung function, biomarkers and risk of asthma. (PMID:18511640; 18848868 )

## Notable Publications

Author	Pubmed ID	Journal	Application
Jinliang Liu	34630847	Oxid Med Cell Longev	WB
Lei Zhao	34582963	Food Chem Toxicol	WB
Kang Wang	36139913	Antioxidants (Basel)	WB


## Storage

Storage:  
Store at -20°C. Stable for one year after shipment.  
Storage Buffer:  
PBS with 0.02% sodium azide and 50% glycerol, pH7.3  
Aliquoting is unnecessary for -20°C storage


For technical support and original validation data for this product please contact:  
T: 4006900926 E: Proteintech-CN@ptglab.com W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.


## Selected Validation Data




Various lysates were subjected to SDS PAGE followed by western blot with 67240-1-Ig (NQO1 antibody) at dilution of 1:20000 incubated at room temperature for 1.5 hours.



Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 67240-1-Ig (NQO1 antibody) at dilution of 1:5000 (under 10x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



Immunohistochemical analysis of paraffin-embedded human colon tissue slide using 67240-1-Ig (NQO1 antibody) at dilution of 1:5000 (under 40x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0).



1X10<sup>6</sup> MCF-7 cells were intracellularly stained with 0.4 ug Anti-Human NQO1 (67240-1-Ig, Clone:1E5G7) and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) at dilution 1:1000 (red), or 0.4 ug Mouse IgG2a Isotype Control (66360-2-Ig, Clone: K11A1B2A2) (blue). Cells were fixed with 4% PFA and permeabilized with Flow Cytometry Perm Buffer (PF00011-C).