For Research Use Only

CDK9 Monoclonal antibody

Catalog Number:67256-1-Ig




Basic Information	Catalog Number: 67256-1-Ig	GenBank Accession Number: BC001968	Purification Method: Protein A purification	
	Size: 1000 µg/ml	GenelD (NCBI): 1025	CloneNo.: 1G6B3	
	Source: Mouse Isotype: IgG1 Immunogen Catalog Number: AG29178	UNIPROT ID: P50750 Full Name: cyclin-dependent kinase 9	Recommended Dilutions: WB 1:5000-1:50000 IF/ICC 1:500-1:2000	
				Calculated MW: 372 aa, 43 kDa
		Observed MW: 38-42 kDa, 55 kDa		
		Applications	Tested Applications: IF/ICC, WB, ELISA Species Specificity: Human, Mouse, Rat	Positive
	WB : HeLa cells, T-47D cells, HCT 116 cells, A431 cells, MOLT-4 cells, TF-1 cells, HSC-T6 cells, ROS1728 cells			
IF/ICC : I	HeLa cells,			
Background Information	CDK9(Cyclin-dependent kinase 9) is a member of the Cdc2-like family of kinases. Its cyclin partners are members of the family of cyclin T (T1, T2a and T2b) and cyclin K. The CDK9/cyclin T complexes appear to be involved in regulating several physiological processes. CDK9 has also been described as the kinase of the TAK complex, which is homologous to the P-TEFb complex and involved in HIV replication. In addition, CDK9 seems to have an anti-apoptotic function in monocytes, that may be related to its control over differentiation of monocytes (PMID: 12432243). CDK9 has two isoforms with the molecular mass of 42 kDa and 55 kDa, and the relative abundance of Cdk9(42kDa) and Cdk9(55kDa) changes in different cell types (PMID: 12706900, 15780980).			
Storage	Storage: Store at -20°C. Stable for one year Storage Buffer: PBS with 0.1% sodium azide and 5 Aliquoting is unnecessary for -20°	50% glycerol pH 7.3.		


 For technical support and original validation data for this product please contact:

 T: 4006900926
 E: Proteintech-CN@ptglab.com
 W: ptgcn.com

This product is exclusively available under Proteintech Group brand and is not available to purchase from any other manufacturer.

Selected Validation Data





CDK9 Merge

Immunofluorescent analysis of (4% PFA) fixed HeLa cells using CDK9 antibody (67256-1-lg, Clone: 1G6B3) at dilution of 1:1000 and Multi-rAbC oraLite Plus 594-Goat Anti-Mouse Recombinant Secondary Antibody (H+L) (RGAM004, 1:500) were used for detection.

Various lysates were subjected to SDS PAGE followed by western blot with 67256-1-1g (CDK9 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.