### For Research Use Only

# ODC1 Monoclonal antibody

Catalog Number:67336-1-lg Featured Product

1 Publications



**Basic Information** 

Catalog Number: 67336-1-lg Size: 1000 ug/ml

Source: Mouse P11926 Full Name: Isotype: lgG1

Immunogen Catalog Number:

AG10927

GenBank Accession Number:

BC025296 GeneID (NCBI): **UNIPROT ID:** 

ornithine decarboxylase 1

Calculated MW: 461 aa, 51 kDa

Observed MW: 51 kDa

**Purification Method:** Protein A purification

CloneNo.:

2A1E3 Recommended Dilutions: WB 1:5000-1:50000 IF/ICC 1:500-1:2000

**Applications** 

**Tested Applications:** WB, IF/ICC, ELISA Cited Applications:

WB

Species Specificity:

human **Cited Species:** mouse

**Positive Controls:** 

WB: MCF-7 cells, HUVEC cells, human placenta tissue

IF/ICC: MCF-7 cells,

## **Background Information**

Ornithine decarboxylase (ODC) is also named as ODC1 and belongs to the Orn/Lys/Arg decarboxylase class-II family. It catalyzes the conversion of ornithine to putrescine, the first step and a major site of regulation of polyamine biosynthesis. The level of ODC is known to be controlled at several sites, namely transcription, translation, and enzyme degradation. Polyamines can stimulate the degradation of ODC as a type of negative feedback control (PMID:8486633). This protein can be phosphorylated in vivo (PMID:8798774). ODC1 can form a homodimer and only the dimer is catalytically active, as the active sites are constructed of residues from both monomers (PMID: 10623504). The molecular mass of ODC1 is 51 kDa, and the homodimer is 106 kDa.

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Tingting Li	38657365	J Pharm Biomed Anal	WB

Storage


Storage:


Store at -20°C. Stable for one year after shipment.


PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage


### **Selected Validation Data**







MCF-7 cells were subjected to SDS PAGE followed by western blot with 67336-1-1g (ODC1 antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours.



Immunofluorescent analysis of (-20°C Methanol) fixed MCF-7 cells using ODC1 antibody (67336-1-Ig, Clone: 2A1E3) at dilution of 1:1000 and CoraLite®488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L) (SA00013-1).