For Research Use Only

TDP-43 (C-terminal) Monoclonal antibody, PBS Only



Catalog Number: 67345-1-PBS

Featured Product

Basic Information

Catalog Number: 67345-1-PBS

Size: 1 mg/ml Source:

Mouse Isotype:

lgG2a

GenBank Accession Number:

NM_007375 Genel D (NCBI): 23435

UNIPROT ID: Q13148 Full Name:

TAR DNA binding protein

Calculated MW: 43 kDa Observed MW: 43-45 kDa Purification Method: Protein A purification

CloneNo.: 1G5F6

Applications

Tested Applications: WB, IF, IHC, ELISA Species Specificity: Human, mouse, rat

Background Information

Transactivation response (TAR), DNA-binding protein of 43 kDa (also known as TARDBP or TDP-43), was first isolated as a transcriptional inactivator binding to the TAR DNA element of the HIV-1 virus. Neumann et al. (2006) found that a hyperphosphorylated, ubiquitinated, and cleaved form of TARDBP, known as pathologic TDP-43, is the major component of the tau-negative and ubiquitin-positive inclusions that characterize amyotrophic lateral sclerosis (ALS) and the most common pathological subtype of frontotemporal lobar degeneration (FTLD-U). 67345-1-Ig is a mouse monoclonal antibody raised against the C-terminal region of human TDP-43.

Storage


Storage:

Store at -80°C.


The product is shipped with ice packs. Upon receipt, store it immediately at -80°C Storage Buffer:

Storage Buffer PBS only

Selected Validation Data




Immunofluorescent analysis of (4% PFA) fixed rat brain tissue using 67345-1-Ig (TDP-43 (C-terminal) antibody) at dilution of 1:500 and CoraLite488-Conjugated AffiniPure Goat Anti-Mouse IgG(H+L). This data was developed using the same antibody clone with 67345-1-PBS in a different storage buffer formulation.



WB result of TDP-43 (C-terminal) antibody (67345-1-lg: 1:30000; incubated at room temperature for 1.5 hours) with sh-Control and sh-TDP-43 (C-terminal) transfected Hela cells. This data was developed using the same antibody clone with 67345-1-PBS in a different storage buffer formulation



Immunohistochemical analysis of paraffinembedded mouse brain tissue slide using 67345-1-Ig (TDP-43 (C-terminal) antibody) at dilution of 1:5000 (under 4x lens). Heat mediated antigen retrieval with Tris-EDTA buffer (pH 9.0). This data was developed using the same antibody clone with 67345-1-PBS in a different storage buffer formulation.



Various lysates were subjected to SDS PAGE followed by western blot with 67345-1-lg (TDP-43 (C-terminal) antibody) at dilution of 1:10000 incubated at room temperature for 1.5 hours. This data was developed using the same antibody clone with 67345-1-PBS in a different storage buffer formulation.