### For Research Use Only

# HDAC9 Monoclonal antibody

Catalog Number:67364-1-lg 2 Publications



**Basic Information** 

Catalog Number: 67364-1-lg

Size: 1700 µg/ml Source: Mouse Isotype:

lgG2a Immunogen Catalog Number:

AG28514

Calculated MW: 1011 aa, 111 kDa Observed MW: 130 kDa

**Purification Method:** 

Protein A purification CloneNo.:

1G6C5 Recommended Dilutions:

WB 1:1000-1:4000

**Applications** 

**Tested Applications:** WB, ELISA

Cited Applications:

WB

Species Specificity:

Human **Cited Species:** human

#### Positive Controls:

WB: HeLa cells, Daudi cells, HepG2 cells, Raji cells, K-562 cells. Ramos cells

## **Background Information**

HDAC9, also named as Histone deacetylase 7B, is a 1011 amino acid protein, which is responsible for the deacetylation of lysine residues on the N-terminal part of the core histones (H2A, H2B, H3 and H4). Histone deacetylation gives a tag for epigenetic repression and plays an important role in transcriptional regulation, cell cycle progression and developmental events. HDAC9 represses MEF2-dependent transcription. HDAC9 is broadly expressed, with highest levels in brain, heart, muscle and testis.

GenBank Accession Number:

BC152405

GeneID (NCBI):

**UNIPROT ID:** 

Q9UKV0

Full Name:

histone deacetylase 9

#### **Notable Publications**

Author	Pubmed ID	Journal	Application
Yingtong Feng	35595735	Cell Death Dis	WB
Jia-Yi Ning	37602334	Int J Ophthalmol	WB

Storage


Store at -20°C. Stable for one year after shipment.

Storage Buffer

PBS with 0.02% sodium azide and 50% glycerol pH 7.3.

Aliquoting is unnecessary for -20°C storage

# Selected Validation Data



Various lysates were subjected to SDS PAGE followed by western blot with 67364-1-1g (HDAC9 antibody) at dilution of 1:2000 incubated at room temperature for 1.5 hours.